arXiv:1704.05829v1 [math.PR] 19 Apr 2017

Kesten's bound for sub-exponential densities on the real line and its multi-dimensional analogues

Dmitri Finkelshtein[1] Pasha Tkachov[2]

April 20, 2017

Abstract

We study sub-exponential probability densities on the real line, which have the property $f^{*n}(s) \sim nf(s)$, $s \to \infty$, and prove Kesten's bound for them, that is $f^{*n}(s) \leq c_\delta(1+\delta)^n f(s)$ for small δ and large s. We introduce a class of regular sub-exponential functions and use it to find an analogue of Kesten's bound for functions on $\mathbb{R}^d$. The results are applied for the study of the fundamental solution to a nonlocal heat-equation.

Keywords: sub-exponential densities, long-tail functions, heavy-tailed distributions, convolution tails, tail-equivalence, asymptotic behavior

2010 Mathematics Subject Classification: 60E05, 45M05, 62E20

1 Introduction

Let F be a probability distribution on $\mathbb{R}$. Denote by $\overline{F}(s) := F\big((s,\infty)\big)$, $s \in \mathbb{R}$ its tail function. For probability distributions F_1, F_2 on $\mathbb{R}$, their convolution $F_1 * F_2$ has the tail function

$$\overline{F_1 * F_2}(s) = \int_{\mathbb{R}} \overline{F}_1(s-\tau)F_2(d\tau) = \int_{\mathbb{R}} \overline{F}_2(s-\tau)F_1(d\tau),$$

where $\overline{F}_1, \overline{F}_2$ are the corresponding tail functions of F_1, F_2.

If a probability distribution F is concentrated on $\mathbb{R}_+ := [0,\infty)$ and $\overline{F}(s) > 0$, $s \in \mathbb{R}$, then, see e.g. [7],

$$\liminf_{s\to\infty} \frac{\overline{F * F}(s)}{\overline{F}(s)} \geq 2. \tag{1.1}$$

If, additionally, F is heavy-tailed, i.e. $\int_{\mathbb{R}} e^{\lambda s}F(ds) = \infty$ for all $\lambda > 0$, then the equality holds in (1.1), see [12]. An important sub-class of heavy-tailed distributions concentrated on $\mathbb{R}_+$ constitute *sub-exponential* ones, for which

$$\lim_{s\to\infty} \frac{\overline{F * F}(s)}{\overline{F}(s)} = 2. \tag{1.2}$$

[1]Department of Mathematics, Swansea University, Singleton Park, Swansea SA2 8PP, U.K. (d.l.finkelshtein@swansea.ac.uk).

[2]Fakultät für Mathematik, Universität Bielefeld, Postfach 110 131, 33501 Bielefeld, Germany (ptkachov@math.uni-bielefeld.de).

Any sub-exponential distribution on $\mathbb{R}_+$ is (right-side) long-tailed on $\mathbb{R}$, see e.g. [7], i.e. (cf. Definition 2.1 below)

$$\lim_{s\to\infty} \frac{\overline{F}(s+t)}{\overline{F}(s)} = 1, \quad \text{for each } t > 0. \tag{1.3}$$

If distributions F_1, F_2 on $\mathbb{R}$ have probability densities $f_1 \geq 0, f_2 \geq 0$, with $\int_{\mathbb{R}} f_1(s)\,ds = \int_{\mathbb{R}} f_2(s)\,ds = 1$, then $F_1 * F_2$ has the density

$$(f_1 * f_2)(s) := \int_{\mathbb{R}} f_1(s-t) f_2(t)\,dt, \quad s \in \mathbb{R}.$$

The density f of a sub-exponential distribution F concentrated on $\mathbb{R}_+$ (i.e. $f(s) = 0$ for $s < 0$) is said to be sub-exponential on $\mathbb{R}_+$ if f is long-tailed, i.e. (1.3) holds with $\overline{F}$ replaced by f (see also Definition 2.1 below), and, cf. (1.2),

$$\lim_{s\to\infty} \frac{(f * f)(s)}{f(s)} = 2. \tag{1.4}$$

It can be shown (see e.g. [2, 13, 14]) that, in this case, for any $n \in \mathbb{N}$,

$$\lim_{s\to\infty} \frac{f^{*n}(s)}{f(s)} = n, \tag{1.5}$$

where $f^{*n} := f * \ldots * f$ ($n-1$ times). Note that, in general, the density of a sub-exponential distribution concentrated on $\mathbb{R}_+$ even being long-tailed does not need to be a sub-exponential one; the corresponding characterisation can be found in [2, 13], see (2.13) below. The property (1.5) implies, in particular, that, for each $\delta > 0$, $n \in \mathbb{N}$, there exists $s_n > 0$, such that $f^{*n}(s) \leq (n+\delta) f(s)$ for $s > s_n$. In many situations, it is important to have similar inequalities 'uniformly' in n, i.e. on a set independent of n. A possible solution is given by the so-called Kesten's bound, see [14, 2]: for a bounded sub-exponential density f on $\mathbb{R}_+$ and for any $\delta > 0$, there exist $s_0 = s_0(\delta)$ and $c_\delta > 0$, such that

$$f^{*n}(s) \leq c_\delta (1+\delta)^n f(s), \quad s > s_0,\ n \in \mathbb{N}. \tag{1.6}$$

For the corresponding results for distributions, see [7, 8, 3, 13]. Kesten's bounds were used to study series of convolutions of distributions on $\mathbb{R}_+$, $\sum_{n=1}^{\infty} \lambda_n F^{*n}$, and of the corresponding densities, $\sum_{n=1}^{\infty} \lambda_n f^{*n}$, appeared in different contexts: starting from the renewal theory (that was the motivation for the original paper [7]) to branching age dependent processes, random walks, queue theory, risk theory and ruin probabilities, compound Poisson processes, and the study of infinitely divisible laws, see e.g. [10, 2, 13, 21, 14, 4] and the references therein.

If F is a probability distribution on the whole $\mathbb{R}$ such that F^+, given by $F^+(B) := F(B \cap \mathbb{R}_+)$ for all Borel $B \subset \mathbb{R}$, is sub-exponential on $\mathbb{R}_+$, then (see e.g. [13, Lemma 3.4]) F is long-tailed on $\mathbb{R}$ and (1.2) holds. The distributions on $\mathbb{R}$ and their densities were considered by several authors, see [20, 19, 17, 21] and others. The reference [21], in particular, gives a review of difficulties appeared in the case of the whole $\mathbb{R}$ and closes several gaps in the preceding results. However, even some basic properties of sub-exponential densities on the whole $\mathbb{R}$ remained open. Namely, in [13, Lemma 4.13], it was shown that if an integrable on $\mathbb{R}$ function f

(i) is right-side long-tail and, being restricted on $\mathbb{R}_+$ and normalized in $L^1(\mathbb{R}_+)$, satisfies (1.4) (we will say then that f is *weekly sub-exponential* on $\mathbb{R}$, cf. Definition 2.8 below), and if

(ii) the condition (2.9) below holds with $b = f$ (in particular, if f decays to 0 at ∞, cf. Definition 2.15),

then (1.4) holds for the original f on $\mathbb{R}$ as well. We generalize this to an analogue of (1.5) for a general $n \in \mathbb{N}$. In particular, we prove in Theorem 2.19 below that

Theorem 1.1. *If an integrable on $\mathbb{R}$ function f is weakly sub-exponential on $\mathbb{R}$ and decays at ∞, then f, being normalized in $L^1(\mathbb{R})$, satisfies* (1.5).

Moreover, in Theorem 2.22, we prove that then (1.6) holds as well. Namely, in a slightly simplified form (cf. Remark 2.23), one has the following result.

Theorem 1.2. *Let f be a weakly sub-exponential probability density on $\mathbb{R}$, such that, for some $B, K, \nu, \rho > 0$,*

$$f(s+\tau) \le K f(s), \quad s > \rho,\ \tau > 0; \qquad f(s) \le \frac{B}{1+|s|^{1+\nu}}, \quad s < -\rho.$$

Then, for each $\delta > 0$, there exist $s_0 = s_0(\delta) > 0$ and $c_\delta > 0$, such that (1.6) *holds.*

Note that the all 'classical' examples of sub-exponential functions satisfy assumptions of Theorem 1.2, see Subsection 3.2.

The multi-dimensional version of the constructions above is much more non-trivial. Currently, there exist at least three different definitions of sub-exponential distributions on $\mathbb{R}^d$ for $d > 1$, see [9, 16, 18]. The variety is mainly related to different possibilities to describe the zones in $\mathbb{R}^d$ where an analogue of the equivalence (1.2) takes place. However, any results about sub-exponential densities in $\mathbb{R}^d$, $d > 1$, seem to be absent at all. In some sense, the properties of the distribution tails and the integrated tails of the corresponding densities are not related in the multi-dimensional case, since, for a probability density a on $\mathbb{R}^d$,

$$1 - \int_{-\infty}^{x_1} \dots \int_{-\infty}^{x_d} a(y)\, dy \neq \int_{x_1}^{\infty} \dots \int_{x_d}^{\infty} a(y)\, dy,$$

unless $d = 1$. Note also that if, e.g. a is radially symmetric, i.e. $a(x) = b(|x|)$, $x \in \mathbb{R}^d$ (here $|x|$ denotes the Euclidean norm on $\mathbb{R}^d$) and b, being normalized, is a sub-exponential density on $\mathbb{R}_+$, then

$$(a * a)(x) := \int_{\mathbb{R}^d} a(x-y)a(y)\, dy = c(|x|), \quad x \in \mathbb{R}^d,$$

for some $c : \mathbb{R}_+ \to \mathbb{R}_+$, i.e. $a * a$ is also radially symmetric, however, asymptotic behaviors of b and c at ∞ are hardly to be compared. Leaving this problem as on open, we concentrate in the present paper on an analogue of Kesten's bound (1.6) in the multi-dimensional case. To do this, we introduce a special class $\widetilde{\mathcal{S}}_{\mathrm{reg},d}$ of regular sub-exponential functions on $\mathbb{R}_+$ (see Definitions 3.1 and 4.11). Functions from this class are either inverse polynomials (i.e. (4.5) holds), or decay at ∞ faster than any polynomial (i.e. (4.15) holds), but slower than any exponential function, with the fastest allowed asymptotic $\exp\big(-\frac{s}{(\log s)^\alpha}\big)$ with $\alpha > 1$, cf. Remark 3.15. Then, in Corollary 4.14, we show the following result.

Theorem 1.3. *Let $a = a(x)$ be a probability density on $\mathbb{R}^d$, such that $a(x) = b(|x|)$, $x \in \mathbb{R}^d$, for some $b \in \widetilde{\mathcal{S}}_{\mathrm{reg},d}$. Then, for each $\delta > 0$ and for each $\alpha < 1$ close enough to 1,*

$$a^{*n}(x) \le c_{\delta,\alpha}(1+\delta)^n a(x)^\alpha, \quad |x| > s_\alpha,\ n \in \mathbb{N} \tag{1.7}$$

for some $c_{\delta,\alpha}, s_\alpha = s_\alpha(\delta) > 0$.

The results of Corollary 4.14 is based on more general Theorem 4.13, which says that if, for some $b \in \widetilde{\mathcal{S}}_{\mathrm{reg},d}$,

$$a(x) \le b^+(|x|), \quad x \in \mathbb{R}^d, \qquad \log b^+(s) \sim \log b(s), \quad s \to \infty,$$

then (1.7) holds with $a(x)$ replaced by $b(|x|)$ in the right hand side.

The paper is organized as follows. In Section 2, we consider properties of general sub-exponential functions on the real line and prove the results which imply Theorems 1.1 and 1.2. In Section 3, we define and study properties of regular sub-exponential functions on the real line and consider the corresponding examples. In Section 4, we prove Theorem 1.3 and its generalizations. Finally, in Appendix, we apply the obtained results to the study of a non-local heat equation.

2 Sub-exponential functions and Kesten's bound on the real line

Definition 2.1. A function $b : \mathbb{R} \to \mathbb{R}_+$ is said to be *(right-side) long-tailed* if there exists $\rho = \rho_b \ge 0$, such that $b(s) > 0$ for all $s \ge \rho$; and, for any $\tau \ge 0$,

$$\lim_{s\to\infty} \frac{b(s+\tau)}{b(s)} = 1. \tag{2.1}$$

Remark 2.2. By [13, formula (2.18)], the convergence in (2.1) is equivalent to the locally uniform in τ convergence, namely, (2.1) can be replaced by the assumption that, for all $h > 0$,

$$\lim_{s\to\infty} \sup_{|\tau|\le h} \left| \frac{b(s+\tau)}{b(s)} - 1 \right| = 0. \tag{2.2}$$

A long-tailed function has to have a 'heavier' tail than any exponential function; namely, the following statement holds.

Lemma 2.3 ([13, Lemma 2.17])**.** *Let $b : \mathbb{R} \to \mathbb{R}_+$ be a long-tailed function. Then, for any $k > 0$,*

$$\lim_{s\to\infty} e^{ks} b(s) = \infty.$$

The constant h in (2.2) may be arbitrary big. It is quite natural to ask what will be if h increases to ∞ consistently with s.

Lemma 2.4 (cf. [13, Lemma 2.19, Proposition 2.20])**.** *Let $b : \mathbb{R} \to \mathbb{R}_+$ be a long-tailed function. Then there exists a function $h : (0,\infty) \to (0,\infty)$, with $h(s) < \frac{s}{2}$ and $\lim\limits_{s\to\infty} h(s) = \infty$, such that, cf. (2.2),*

$$\lim_{s\to\infty} \sup_{|\tau|\le h(s)} \left| \frac{b(s+\tau)}{b(s)} - 1 \right| = 0. \tag{2.3}$$

Following [13], we will say then that b is h-insensitive. Of course, for a given long-tailed function b, the function h that fulfills (2.3) is not unique, see also [13, Proposition 2.20].

The convergence in (2.1) will not be, in general, monotone in s. To get this monotonicity, we consider the following class of functions.

Definition 2.5. A function $b : \mathbb{R} \to \mathbb{R}_+$ is said to be *(right-side) tail-log-convex*, if there exists $\rho = \rho_b > 0$ such that $b(s) > 0$, $s \geq \rho$, and the function $\log b$ is convex on $[\rho, \infty)$.

Remark 2.6. It is well-known that any function which is convex on an open interval is continuous there. Therefore, a tail-log-convex function $b = \exp(\log b)$ is continuous on (ρ_b, ∞) as well.

Lemma 2.7. *Let $b : \mathbb{R} \to \mathbb{R}_+$ be tail-log-convex, with $\rho = \rho_b$. Then, for any $\tau > 0$, the function $\dfrac{b(s+\tau)}{b(s)}$ is non-decreasing in $s \in [\rho, \infty)$.*

Proof. Take any $s_1 > s_2 \geq \rho$. Set $B(s) := \log b(s) \leq 0$, $s \in [\rho, \infty)$. Then the desired inequality

$$\frac{b(s_1+\tau)}{b(s_1)} \geq \frac{b(s_2+\tau)}{b(s_2)}$$

is equivalent to

$$B(s_1+\tau) + B(s_2) \geq B(s_2+\tau) + B(s_1).$$

Since B is convex, we have, for $\lambda = \dfrac{\tau}{s_1 - s_2 + \tau} \in (0,1)$,

$$\begin{aligned} B(s_1) = B\big(\lambda s_2 + (1-\lambda)(s_1+\tau)\big) &\leq \lambda B(s_2) + (1-\lambda) B(s_1+\tau), \\ B(s_2+\tau) = B\big((1-\lambda)s_2 + \lambda(s_1+\tau)\big) &\leq (1-\lambda) B(s_2) + \lambda B(s_1+\tau), \end{aligned}$$

that implies the needed inequality. □

Because of the terminology mentioned in the introduction, we will use the following definition.

Definition 2.8. We will say that a function $b : \mathbb{R} \to \mathbb{R}_+$ is *weakly (right-side) sub-exponential on* $\mathbb{R}$ if b is long-tailed, $b \in L^1(\mathbb{R}_+)$, and the function

$$b_+(s) := \mathbb{1}_{\mathbb{R}_+}(s) \left(\int_{\mathbb{R}_+} b(\tau) d\tau \right)^{-1} b(s), \quad s \in \mathbb{R}, \tag{2.4}$$

satisfies the following asymptotic relation

$$\begin{aligned} (b_+ * b_+)(s) &= \int_{\mathbb{R}} b_+(s-\tau) b_+(\tau)\, d\tau \\ &= \int_0^s b_+(s-\tau) b_+(\tau)\, d\tau \sim 2 b_+(s), \quad s \to \infty. \end{aligned} \tag{2.5}$$

The next statement shows that a long-tailed tail-log-convex function is weakly sub-exponential on $\mathbb{R}$ provided that it decays at ∞ fast enough.

Lemma 2.9 (cf. [13, Theorem 4.15]). *Let $b : \mathbb{R} \to \mathbb{R}_+$ be a long-tailed tail-log-convex function such that $b \in L^1(\mathbb{R}_+)$. Suppose that, for a function $h : (0,\infty) \to (0,\infty)$, with $h(s) < \frac{s}{2}$ and $\lim\limits_{s\to\infty} h(s) = \infty$, the asymptotic (2.3) holds, and that*

$$\lim_{s\to\infty} s\, b\big(h(s)\big) = 0. \tag{2.6}$$

Then b is weakly sub-exponential on $\mathbb{R}$.

Remark 2.10. Let $b : \mathbb{R} \to \mathbb{R}_+$ be a weakly sub-exponential function on $\mathbb{R}$. Then, by (2.4), (2.5), we have

$$\int_0^s b(s-\tau)b(\tau)\,d\tau \sim 2\bigg(\int_{\mathbb{R}_+} b(\tau)d\tau\bigg)b(s), \quad s\to\infty. \tag{2.7}$$

Definition 2.11. We will say that a function $b : \mathbb{R} \to \mathbb{R}_+$ is *strongly (right-side) sub-exponential on* $\mathbb{R}$ if b is long-tailed, $b \in L^1(\mathbb{R})$, and the following asymptotic relation holds, cf. (2.5), (2.7),

$$(b * b)(s) = \int_{\mathbb{R}} b(s-\tau)b(\tau)\,d\tau \sim 2\bigg(\int_{\mathbb{R}} b(\tau)d\tau\bigg)b(s), \quad s\to\infty. \tag{2.8}$$

Remark 2.12. By [13, Lemma 4.12], a strongly sub-exponential function on $\mathbb{R}$ is weakly sub-exponential there. The following lemma presents a sufficient condition to get the converse.

Lemma 2.13 (cf. [13, Lemma 4.13]). *Let $b \in L^1(\mathbb{R} \to \mathbb{R}_+)$ be a weakly sub-exponential function on $\mathbb{R}$. Suppose that there exists $\rho = \rho_b > 0$ and $K = K_b > 0$ such that*

$$b(s+\tau) \le Kb(s), \quad s > \rho,\ \tau > 0. \tag{2.9}$$

Then (2.8) holds, i.e. b is strongly sub-exponential on $\mathbb{R}$.

Remark 2.14. For $b \in L^1(\mathbb{R} \to \mathbb{R}_+)$, condition (2.9) yields that $\sup_{t\ge s} b(t) \to 0$, $s \to \infty$. In particular $b(s) \to 0$, as $s \to \infty$.

An evident sufficient condition which ensures (2.9) is that b is decreasing on $[\rho,\infty)$. Consider the corresponding definition.

Definition 2.15. A function $b : \mathbb{R} \to \mathbb{R}_+$ is said to be *(right-side) tail-decreasing* if there exists a number $\rho = \rho_b \ge 0$ such that $b = b(s)$ is strictly decreasing on $[\rho,\infty)$ to 0. In particular, $b(s) > 0$, $s \ge \rho$.

Proposition 2.16. *Let $b : \mathbb{R} \to \mathbb{R}_+$ be a tail-decreasing function. Let $h : (0,\infty) \to (0,\infty)$, with $h(s) < \frac{s}{2}$ and $\lim\limits_{s\to\infty} h(s) = \infty$. Then (2.3) holds, if and only if*

$$\lim_{s\to\infty} \frac{b(s \pm h(s))}{b(s)} = 1. \tag{2.10}$$

Proof. Let $\rho = \rho_b > 0$ be as in the Definition 2.15. Then, for the given h and for any $s > 2\rho$, one has that $s - h(s) > \frac{s}{2} > \rho$. Hence, for a fixed $s > 2\rho$,

the function $b(s+\tau)$ is decreasing in $\tau \in [-h(s), h(s)]$. Therefore, considering separately positive and negative τ, one gets that, for all $s > 2\rho$,

$$\sup_{|\tau|\le h(s)} \left| \frac{b(s+\tau)}{b(s)} - 1 \right| = \max\left\{ 1 - \frac{b(s+h(s))}{b(s)}, \frac{b(s-h(s))}{b(s)} - 1 \right\},$$

that yields the statement. □

Proposition 2.17. *Let $b : \mathbb{R} \to \mathbb{R}_+$ be a weakly sub-exponential function on $\mathbb{R}$, such that (2.9) holds. Let $b_1, b_2 \in L^1(\mathbb{R} \to \mathbb{R}_+)$ and there exist constants $c_1, c_2 \ge 0$, such that*

$$\lim_{s\to\infty} \frac{b_j(s)}{b(s)} = c_j, \quad j = 1, 2. \tag{2.11}$$

Then

$$\lim_{s\to\infty} \frac{(b_1 * b_2)(s)}{b(s)} = c_1 \int_{\mathbb{R}} b_2(\tau)\, d\tau + c_2 \int_{\mathbb{R}} b_1(\tau)\, d\tau. \tag{2.12}$$

Proof. We will follow ideas of the proofs for [13, Lemma 4.13, Lemma 4.9]. Let b_+ be given by (2.4). Since evidently, cf. (2.1), b_+ is also a long-tailed function, and (2.5) holds, we have, by [13, Theorem 4.7], that, for any $h : (0,\infty) \to (0,\infty)$, with $h(s) < \frac{s}{2}$ and $\lim_{s\to\infty} h(s) = \infty$, the following asymptotic holds

$$\int_{h(s)}^{s-h(s)} b_+(s-\tau) b_+(\tau)\, d\tau = o(b_+(s)), \quad s \to \infty. \tag{2.13}$$

In particular, one can choose and fix an increasing function h, such that the properties above hold. Note that $h(s) \le \tau \le s - h(s)$ implies $\tau > 0$ and $s - \tau \ge h(s) > 0$, hence b_+ may be replaced by b in (2.13).

Next, for any $b_1, b_2 \in L^1(\mathbb{R} \to \mathbb{R}_+)$, one can easily get that

$$\begin{aligned}(b_1 * b_2)(s) = &\int_{-\infty}^{-h(s)} \big(b_1(s-\tau) b_2(\tau) + b_1(\tau) b_2(s-\tau)\big)\, d\tau \\ &+ \int_{-h(s)}^{h(s)} \big(b_1(s-\tau) b_2(\tau) + b_1(\tau) b_2(s-\tau)\big)\, d\tau \\ &+ \int_{h(s)}^{s-h(s)} b_1(s-\tau) b_2(\tau)\, d\tau.\end{aligned}$$

Take an arbitrary $\delta \in (0,1)$. By (2.11), (2.3), and (2.13) (the latter, with b_+ replaced by b), there exist $K > 0$ and $\rho > 0$, such that (2.9) holds and, for all $s \ge h(\rho)$,

$$\big|b_j(s) - c_j b(s)\big| \le \delta b(s), \tag{2.14}$$

$$\sup_{|\tau|\le h(s)} \big|b(s+\tau) - b(s)\big| \le \delta b(s), \tag{2.15}$$

$$\int_{h(s)}^{s-h(s)} b(s-\tau) b(\tau)\, d\tau \le \delta b(s), \tag{2.16}$$

$$\int_{-\infty}^{-h(s)} b_j(\tau) d\tau + \int_{h(s)}^{\infty} b_j(\tau) d\tau \le \delta, \quad j = 1, 2. \tag{2.17}$$

For any $s \geq \rho > h(\rho)$, we trivially have that $\tau \leq -h(s) < 0$ implies, $s - \tau > s > h(\rho)$. Then, by (2.14),

$$b_j(s-\tau) \leq (c_j + \delta) b(s-\tau), \quad \tau < -h(s),\ j = 1, 2;$$

and, therefore, by (2.9) and (2.17), for $s \geq \rho > h(\rho)$,

$$\begin{aligned}
&\int_{-\infty}^{-h(s)} \big(b_1(s-\tau)b_2(\tau) + b_1(\tau)b_2(s-\tau)\big)\, d\tau \\
&\leq \int_{-\infty}^{-h(s)} \big(K(c_1+\delta)b(s)b_2(\tau) + K(c_2+\delta)b_1(\tau)b(s)\big)\, d\tau \\
&\leq \delta K(c_1 + c_2 + 2\delta) b(s).
\end{aligned}$$

Next, $-h(s) \leq \tau \leq h(s)$ implies $s - \tau \geq s - h(s) > h(s) \geq h(\rho)$ for $s \geq \rho$ (recall that h is increasing). Then, by (2.14), (2.15),

$$\begin{aligned}
&\left| \int_{-h(s)}^{h(s)} b_1(s-\tau)b_2(\tau)\, d\tau - c_1 b(s) \int_{-h(s)}^{h(s)} b_2(\tau)\, d\tau \right| \\
&\leq \int_{-h(s)}^{h(s)} \big| b_1(s-\tau) - c_1 b(s-\tau) \big| b_2(\tau)\, d\tau \\
&\quad + c_1 \int_{-h(s)}^{h(s)} \big| b(s-\tau) - b(s) \big| b_2(\tau)\, d\tau \leq \delta B_2 (1 + c_1) b(s).
\end{aligned}$$

Next, $h(s) \leq \tau \leq s - h(s)$ implies $\min\{s-\tau, \tau\} \geq h(s) \geq h(\rho)$ for $s \geq \rho > h(\rho)$; and hence we have, by (2.14), (2.16),

$$\begin{aligned}
\int_{h(s)}^{s-h(s)} b_1(s-\tau)b_2(\tau)\, d\tau &\leq (c_1+\delta)(c_2+\delta) \int_{h(s)}^{s-h(s)} b(s-\tau)b(\tau)\, d\tau \\
&\leq \delta (c_1+\delta)(c_2+\delta) b(s).
\end{aligned}$$

Set $B_j := \int_{\mathbb{R}} b_j(s)\, ds$. The evident identity

$$B_j = \int_{-h(s)}^{h(s)} b_j(\tau)\, d\tau + \int_{-\infty}^{-h(s)} b_j(\tau) d\tau + \int_{h(s)}^{\infty} b_j(\tau) d\tau, \quad j = 1, 2,$$

implies, for $s \geq \rho$,

$$\big| (b_1 * b_2)(s) - (c_1 B_2 + c_2 B_1) b(s) \big| \leq \delta M b(s),$$

where

$$M := B_2(1 + c_1) + B_1(1 + c_2) + c_1 + c_2 + K(c_1 + c_2 + 2\delta) + (c_1 + \delta)(c_2 + \delta).$$

This yields the desired convergence (2.12). □

Corollary 2.18. *The property of an integrable on $\mathbb{R}$ function to be weekly sub-exponential on $\mathbb{R}$ depends on its behaviour at $+\infty$ only. Namely, for a weakly sub-exponential on $\mathbb{R}$ function $b \in L^1(\mathbb{R} \to \mathbb{R}_+)$ and for any $c \in L^1(\mathbb{R} \to \mathbb{R}_+)$ and $s_0 \in \mathbb{R}$, the function $\tilde{b}(s) = \mathbb{1}_{(-\infty, s_0)}(s) c(s) + \mathbb{1}_{[s_0, \infty)}(s) b(s)$ is weakly sub-exponential on $\mathbb{R}$, cf. also Theorem 3.5 below.*

Now one gets a generalization of Lemma 2.13.

Theorem 2.19. *Let $b \in L^1(\mathbb{R} \to \mathbb{R}_+)$ be a weakly sub-exponential on $\mathbb{R}$ function, such that (2.9) holds (for example, let b be tail-decreasing). Then*

$$\lim_{s\to\infty} \frac{b^{*n}(s)}{b(s)} = n\Big(\int_{\mathbb{R}} b(\tau)\,d\tau\Big)^{n-1}, \quad n \geq 2, \tag{2.18}$$

where $b^{*n}(s) = (\underbrace{b * \ldots * b}_{n})(s)$, $s \in \mathbb{R}$.

Proof. Take in Proposition 2.17, $b_1 = b_2 = b$, i.e. $c_1 = c_2 = 1$. Then, for $B := \int_{\mathbb{R}} b(\tau)\,d\tau$, one gets $b^{*2}(s) \sim 2Bb(s)$, $s \to \infty$. Proving by induction, assume that $b^{*(n-1)}(s) \sim (n-1)B^{n-2}b(s)$, $s \to \infty$, $n \geq 3$. Take in Proposition 2.17, $b_1 = b$, $b_2 = b^{*(n-1)}$, $c_1 = 1$, $c_2 = (n-1)B^{n-2}$, then, since $\int_{\mathbb{R}} b^{*(n-1)}(\tau)\,d\tau = B^{n-1}$, one gets

$$\lim_{s\to\infty} \frac{b^{*n}(s)}{b(s)} = B^{n-1} + B(n-1)B^{(n-2)} = nB^{(n-1)}. \qquad \square$$

Consider now some general statements on the Euclidean space $\mathbb{R}^d$, $d \in \mathbb{N}$. Fix the Borel σ-algebra $\mathcal{B}(\mathbb{R}^d)$ there. All functions on $\mathbb{R}^d$ in the sequel are supposed to be $\mathcal{B}(\mathbb{R}^d)$-measurable. Let $0 \leq a \in L^1(\mathbb{R}^d)$ be a fixed probability density on $\mathbb{R}^d$, i.e.

$$\int_{\mathbb{R}^d} a(x)\,dx = 1. \tag{2.19}$$

Let $f : \mathbb{R} \to \mathbb{R}$; we will say that the convolution

$$(a * f)(x) := \int_{\mathbb{R}^d} a(x-y)f(y)\,dy, \quad x \in \mathbb{R}^d$$

is well-defined if the function $y \mapsto a(x-y)f(y)$ belongs to $L^1(\mathbb{R}^d)$ for a.a. $x \in \mathbb{R}^d$. In particular, this holds if $f \in L^\infty(\mathbb{R}^d)$. Next, for a function $\phi : \mathbb{R}^d \to (0, +\infty)$, we define, for any $f : \mathbb{R}^d \to \mathbb{R}$,

$$\|f\|_\phi := \sup_{x\in\mathbb{R}^d} \frac{|f(x)|}{\phi(x)} \in [0, \infty].$$

Proposition 2.20. *Let a function $\phi : \mathbb{R}^d \to (0, +\infty)$ be such that $a * \phi$ is well-defined, $\|a\|_\phi < \infty$, and, for some $\gamma \in (0, \infty)$,*

$$\frac{(a * \phi)(x)}{\phi(x)} \leq \gamma, \quad x \in \mathbb{R}^d. \tag{2.20}$$

Then

$$a^{*n}(x) \leq \gamma^{n-1}\|a\|_\phi \phi(x), \quad x \in \mathbb{R}^d.$$

Proof. For any $f : \mathbb{R}^d \to \mathbb{R}$, with $\|f\|_\phi < \infty$, we have, for $x \in \mathbb{R}^d$,

$$\left|\frac{(a * f)(x)}{\phi(x)}\right| \leq \int_{\mathbb{R}^d} \frac{a(y)\phi(x-y)}{\phi(x)} \frac{|f(x-y)|}{\phi(x-y)} dy \leq \frac{a * \phi(x)}{\phi(x)} \|f\|_\phi \leq \gamma\|f\|_\phi.$$

In particular, since $\|a\|_\phi < \infty$, one gets

$$\|a * a\|_\phi \leq \gamma\|a\|_\phi < \infty.$$

Proceeding inductively, one gets

$$\|a^{*n}\|_\phi \le \gamma \|a * a^{n-1}\|_\phi \le \gamma^{n-1}\|a\|_\phi < \infty,$$

that yields the statement. □

Proposition 2.21. *Let a function $\omega : \mathbb{R}^d \to (0, +\infty)$ be such that, for any $\lambda > 0$, the set*

$$\Omega_\lambda := \Omega_\lambda(\omega) := \big\{x \in \mathbb{R}^d : \omega(x) < \lambda\big\} \neq \emptyset. \tag{2.21}$$

Suppose further

$$\eta := \limsup_{\lambda \to 0+} \sup_{x \in \Omega_\lambda} \frac{(a * \omega)(x)}{\omega(x)} \in (0, \infty). \tag{2.22}$$

Then, for any $\delta \in (0,1)$, there exists $\lambda = \lambda(\delta, \omega) \in (0,1)$, such that (2.20) *holds, with*

$$\phi(x) := \omega_\lambda(x) := \min\big\{\lambda, \omega(x)\big\}, \quad x \in \mathbb{R}^d, \tag{2.23}$$

and $\gamma := \max\{1, (1+\delta)\eta\}$.

Proof. By (2.23), for an arbitrary $\lambda > 0$, we have $\omega_\lambda(x) \le \lambda$, $x \in \mathbb{R}^d$; then $(a * \omega_\lambda)(x) \le \lambda$, $x \in \mathbb{R}^d$, as well. In particular, cf. (2.23),

$$(a * \omega_\lambda)(x) \le \omega_\lambda(x), \quad x \in \mathbb{R}^d \setminus \Omega_\lambda. \tag{2.24}$$

Next, by (2.22), for any $\delta > 0$ there exists $\lambda = \lambda(\delta) \in (0,1)$ such that

$$\sup_{x \in \Omega_\lambda} \frac{(a * \omega)(x)}{\omega(x)} - \eta \le \delta\eta,$$

in particular,

$$(a * \omega)(x) \le (1+\delta)\eta\omega(x) = (1+\delta)\eta\omega_\lambda(x), \quad x \in \Omega_\lambda.$$

Therefore, for all $x \in \Omega_\lambda$,

$$(a * \omega_\lambda)(x) = (a * \omega)(x) - \big(a * (\omega - \omega_\lambda)\big)(x) \le (1+\delta)\eta\omega_\lambda(x), \tag{2.25}$$

where we used the obvious inequality: $\omega \ge \omega_\lambda$. By (2.24) and (2.25), one gets the statement. □

We are ready to prove now Kesten's bound on $\mathbb{R}$.

Theorem 2.22. *Let $b \in L^1(\mathbb{R} \to \mathbb{R}_+)$ be a bounded weakly sub-exponential on $\mathbb{R}$ function with $\int_{\mathbb{R}} b(s)\,ds = 1$, such that* (2.9) *holds. Suppose that there exists a bounded weakly sub-exponential function $c \in L^1(\mathbb{R} \to \mathbb{R}_+)$, such that* (2.9) *holds with b replaced by c, and, for some $D > 0$ and $s' > 0$,*

$$b(-s) \le D\,c(s), \quad s > s'. \tag{2.26}$$

Then, for any $\delta \in (0,1)$, there exist $C_\delta > 0$ and $s_0 = s_0(\delta) > 0$, such that

$$b^{*n}(s) \le C_\delta (1+\delta)^n b(s), \quad s > s_0, n \in \mathbb{N}. \tag{2.27}$$

Remark 2.23. Because of the examples in Section 3 below, to get (2.26), it is enough to assume that, for some $B, \nu > 0$,

$$b(s) \le \frac{B}{1+|s|^{1+\nu}}, \quad s < -s'.$$

Proof. We fix $\delta \in (0,1)$. By Lemma 2.3, for any $\varepsilon_1 \in (0,1)$, such that $(1+\varepsilon_1) \le (1+\delta)^{\frac{1}{4}}$, and for fixed $C, k > 0$, there exists $s_1 > s'$, such that

$$\begin{aligned} &\min\{b(s), c(s)\} \ge Ce^{-ks}, \qquad s \ge s_1, \\ &\int_{s_1}^{\infty} \big(b(s)+c(s)\big)ds \le \frac{\varepsilon_1}{2}, \end{aligned} \tag{2.28}$$

and the inequality in (2.26) holds for $s > s_1$. We fix $\lambda_1 \le \frac{\varepsilon_1}{4s_1}$. Then we have

$$\int_{-s_1}^{s_1} \big(\max\{\lambda_1, b(s)\} - b(s)\big)ds \le 2s_1\lambda_1 \le \frac{\varepsilon_1}{2}. \tag{2.29}$$

We define

$$\tilde{b}(s) := \mathbb{1}_{(-\infty,-s_1)}(s)c(-s) + \mathbb{1}_{[-s_1,s_1]}(s)\max\{\lambda_1, b(s)\} + \mathbb{1}_{(s_1,\infty)}(s)b(s).$$

In the sequel, let $\|\cdot\|_1$ denote the norm in $L^1(\mathbb{R})$. By (2.28), (2.29), $\|\tilde{b}-b\|_1 \le \varepsilon_1$. Therefore,

$$\|\tilde{b}\|_1 \le 1+\varepsilon_1. \tag{2.30}$$

By Corollary 2.18, both functions $s \mapsto \tilde{b}(s)$ and $s \mapsto \tilde{b}(-s)$ are weakly subexponential on $\mathbb{R}$. Hence by Theorem 2.19, for any $k, m \in \mathbb{N}$,

$$\lim_{s\to\pm\infty} \frac{\tilde{b}^{*m}(s)}{\tilde{b}^{*k}(s)} = \lim_{s\to\pm\infty} \frac{m\|\tilde{b}\|_1^{m-1}\tilde{b}(s)}{k\|\tilde{b}\|_1^{k-1}\tilde{b}(s)} = \frac{m}{k}\|\tilde{b}\|_1^{m-k}.$$

In particular, for any $\varepsilon_2 \in (0,1)$, such that $(1+\varepsilon_2) \le (1+\delta)^{\frac{1}{4}}$, there exist $m_0 \in \mathbb{N}$ and $s_2 \ge s_1$, such that, for $\omega := \tilde{b}^{*m_0}$,

$$\frac{\tilde{b}*\omega(s)}{\omega(s)} \le \|\tilde{b}\|_1(1+\varepsilon_2), \qquad |s| \ge s_2, \tag{2.31}$$

$$m_0\|\tilde{b}\|_1^{m_0-1}(1-\varepsilon_2) \le \frac{\omega(s)}{\tilde{b}(s)} \le m_0(1+\varepsilon_2)\|\tilde{b}\|_1^{m_0-1}, \qquad |s| \ge s_2. \tag{2.32}$$

For any $r > 0$, $\operatorname*{ess\,inf}_{s\in[-r,r]} \tilde{b}(s) > 0$. It is straightforward to check then by induction, that

$$\operatorname*{ess\,inf}_{-r\le s\le r} \omega(s) = \operatorname*{ess\,inf}_{-r\le s\le r} \tilde{b}^{*m_0}(s) > 0, \qquad r > 0. \tag{2.33}$$

On the other hand, by Remark 2.14 and (2.32), we have

$$\lim_{s\to\pm\infty} \omega(s) = \lim_{s\to\pm\infty} \tilde{b}(s) = 0. \tag{2.34}$$

Hence there exists $\lambda_2 \in (0, \lambda_1]$, such that for any $\lambda \in (0, \lambda_2)$, the set Ω_λ defined by (2.21) will be a non-empty subset of $(-\infty, -s_\lambda) \cup (s_\lambda, \infty)$, where $s_\lambda \to \infty$, as $\lambda \searrow 0$. Therefore, by (2.31), the condition (2.22) holds with

$$a(s) = \frac{\tilde{b}(s)}{\|\tilde{b}\|_1}, \quad s \in \mathbb{R}^1, \qquad \eta = (1+\varepsilon_1)(1+\varepsilon_2).$$

Then, by Proposition 2.21, for any $\varepsilon_3 \in (0,1)$ such that $(1+\varepsilon_3) \le (1+\delta)^{\frac{1}{4}}$, there exists $\lambda_3 \in (0, \min\{\lambda_2, 1\})$, such that (2.20) holds with

$$\phi(s) = \min\{\lambda_3, \omega(s)\}, \qquad \gamma = (1+\varepsilon_1)(1+\varepsilon_2)(1+\varepsilon_3), \tag{2.35}$$

and a given above. By (2.32), (2.33) and since $\tilde{b}$ is bounded, we have $\|\tilde{b}\|_\phi < \infty$. Hence, by Proposition 2.20,

$$\tilde{b}^{*n}(s) \le \gamma^{n-1} \|\tilde{b}\|_1^{n-1} \|\tilde{b}\|_\phi \min\{\lambda_3, \omega(s)\}, \qquad s \in \mathbb{R},\ n \in \mathbb{N}.$$

By (2.34), (2.32), there exists $s_0 \ge s_2 \ge s_1$, such that

$$\min\{\lambda_3, \omega(s)\} \le C_1 \tilde{b}(s) = C_1 b(s), \quad s > s_0$$

for some $C_1 = C_1(m_0, \delta) > 0$. By (2.35) and (2.30), $\gamma \|\tilde{b}\|_1 \le 1 + \delta$. Since, by (2.26) and the construction of $\tilde{b}$, $b(s) \le \tilde{b}(s)$, $s \in \mathbb{R}$, one gets that $b^{*n}(s) \le \tilde{b}^{*n}(s)$ for $s \in \mathbb{R}$, $n \in \mathbb{N}$. As a result, (2.27) holds with $C_\delta = \frac{DC_1}{1+\delta}$. □

3 Regular sub-exponential densities on $\mathbb{R}$

We are going to obtain an analogue of Kesten's bound on $\mathbb{R}^d$ with $d > 1$, at least for radially symmetric functions. Our technique will require to deal with functions $b(|x|)^\alpha$, $x \in \mathbb{R}^d$, where b is a sub-exponential function on $\mathbb{R}_+$ and $\alpha < 1$ is close enough to 1; in particular, we have to be sure that b^α is also sub-exponential on $\mathbb{R}_+$. Moreover, to weaken the condition of radial symmetry, we will allow double-side estimates by functions of the form $p(|x|)b(|x|)$ for appropriate p on $\mathbb{R}_+$ (say, polynomial). Again, we will need have to check whether the functions pb is also sub-exponential on $\mathbb{R}_+$. To check such a stability of the class of sub-exponential on $\mathbb{R}_+$ functions with respect to power and multiplicative perturbations, we have to reduce the class to appropriately regular sub-exponential functions. Then the mentioned stability takes place, see Theorem 3.5 and Proposition 3.11. The examples of regular sub-exponential functions are given in Subsection 3.2. The analogues of Kesten's bound on $\mathbb{R}^d$ are considered in Section 4.

3.1 Main properties

Definition 3.1. Let $\mathcal{S}_{\mathrm{reg}}$ be the set of all functions $b : \mathbb{R} \to \mathbb{R}_+$ such that

1) $b \in L^1(\mathbb{R}_+)$ and b is bounded on $\mathbb{R}$;

2) there exists $\rho = \rho_b > 1$, such that b is log-convex and strictly decreasing to 0 on $[\rho, \infty)$ (i.e. b is simultaneously tail-decreasing and tail-log-convex), and (without loss of generality) $b(\rho) \le 1$;

3) there exist $\delta = \delta_b \in (0,1)$ and an increasing function $h = h_b : (0,\infty) \to (0,\infty)$, with $h(s) < \dfrac{s}{2}$ and $\lim\limits_{s\to\infty} h(s) = \infty$, such that the asymptotic (2.10) holds, and, cf. (2.6),

$$\lim_{s\to\infty} b\big(h(s)\big)s^{1+\delta} = 0. \tag{3.1}$$

For any $n \in \mathbb{N}$, we denote by $\mathcal{S}_{\mathrm{reg},n}$ the subclass of functions b from $\mathcal{S}_{\mathrm{reg}}$ such that

$$\int_{-\infty}^{\rho} b(s)\,ds + \int_{\rho}^{\infty} b(s)s^{n-1}\,ds < \infty. \tag{3.2}$$

Remark 3.2. It is worth noting again that, for a tail-decreasing function, (2.10) implies that b is long-tailed.

Remark 3.3. By Lemma 2.9, any function $b \in \mathcal{S}_{\mathrm{reg}}$ is weakly sub-exponential on $\mathbb{R}$. Moreover, by Lemma 2.13, any function $b \in \mathcal{S}_{\mathrm{reg},1}$ is strongly sub-exponential on $\mathbb{R}$.

Remark 3.4. Let $b \in \mathcal{S}_{\mathrm{reg}}$, and $s_0 > 0$ be such that $h(2s_0) > \rho$. Then the monotonicity of b and h implies $b(s) \le b(h(2s))$, $s > s_0$; and hence, because of (3.1), for $B := 2^{-1-\delta}$, there exists $s_1 \ge s_0$, such that

$$b(s) \le \frac{B}{s^{1+\delta}}, \quad s \ge s_1. \tag{3.3}$$

Below we will show that $\mathcal{S}_{\mathrm{reg}}$ and $\mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$ are closed under some simple transformations of functions. For an arbitrary function $b \in \mathcal{S}_{\mathrm{reg}}$, we consider the following transformed functions:

1) for fixed $p > 0$, $q > 0$, $r \in \mathbb{R}$, we set

$$\widetilde{b}(s) := pb(qs + r), \quad s \in \mathbb{R}; \tag{3.4}$$

2) for a fixed $s_0 > 0$ and a fixed bounded function $c : \mathbb{R} \to \mathbb{R}_+$, we set

$$\breve{b}(s) := \mathbb{1}_{(-\infty,s_0)}(s)c(s) + \mathbb{1}_{[s_0,\infty)}(s)b(s), \quad s \in \mathbb{R}; \tag{3.5}$$

3) for any $\alpha \in (0,1]$, we denote

$$b_\alpha(s) := \big(b(s)\big)^\alpha, \quad s \in \mathbb{R}.$$

Theorem 3.5. *1) Let $b \in \mathcal{S}_{\mathrm{reg}}$. Then the functions $\widetilde{b}$ and $\breve{b}$ defined in* (3.4) *and* (3.5)*, correspondingly, also belong to $\mathcal{S}_{\mathrm{reg}}$ for all admissible values of their parameters. If, additionally, there exists $\alpha' \in (0,1)$ such that $b_{\alpha'} \in L^1(\mathbb{R}_+)$, then there exists $\alpha_0 \in (\alpha',1)$, such that $b_\alpha \in \mathcal{S}_{\mathrm{reg}}$ for all $\alpha \in [\alpha_0,1]$.*

2) Let $b \in \mathcal{S}_{\mathrm{reg},n}$ for some $n \in \mathbb{N}$. Then $\widetilde{b} \in \mathcal{S}_{\mathrm{reg},n}$. If, additionally, the function c in (3.5) *is integrable on $(-\infty, s_0)$, then $\breve{b} \in \mathcal{S}_{\mathrm{reg},n}$. Finally, if there exists $\alpha' \in (0,1)$ such that* (3.2) *holds for $b = b_{\alpha'}$, then there exists $\alpha_0 \in (\alpha',1)$, such that $b_\alpha \in \mathcal{S}_{\mathrm{reg},n}$ for all $\alpha \in [\alpha_0,1]$. Moreover, in the latter case, there exist $B_0 > 0$ and $\rho_0 > 0$, such that, for all $\alpha \in (\alpha_0,1]$,*

$$\int_{\mathbb{R}} \big(b(s-\tau)\big)^\alpha \big(b(\tau)\big)^\alpha\,d\tau \le B_0\big(b(s)\big)^\alpha, \quad s \ge \rho_0, \tag{3.6}$$

Proof. It is very straightforward to check that if b is long-tailed, tail-decreasing and tail-log-convex, then $\widetilde{b}, \breve{b}, b_\alpha$ also have these properties for all admissible values of their parameters. Let $h : (0,\infty) \to (0,\infty)$ be such that $h(s) < \dfrac{s}{2}$, $\lim\limits_{s\to\infty} h(s) = \infty$, and (2.10) hold. Let also (3.1) hold for some $\delta > 0$.

(i) Evidently, both (2.10) and (3.1) hold, with b replaced by $\breve{b}$. Next, $\breve{b} \in L^1(\mathbb{R}_+)$ and $\breve{b}$ is bounded. Hence $\breve{b} \in \mathcal{S}_{\rm reg}$. If $b \in \mathcal{S}_{{\rm reg},n}$ and c is integrable on $(-\infty, s_0)$, then (3.2) holds for b replaced by $\breve{b}$. Cf. also Corollary 2.18.

(ii) Set, for the given $q > 0$, $r \in \mathbb{R}$,

$$\widetilde{h}(s) := \frac{1}{q} h(qs + r) - \frac{r}{2q} \mathbb{1}_{\mathbb{R}_+}(r), \quad s \in [s_1, \infty),$$

where $s_1 > 0$ is such that $qs_1 + r > 0$ and $h(qs + r) > \frac{r}{2q}$ for all $s \ge s_1$. Clearly, $\widetilde{h}$ is increasing on $[s_1, \infty)$, $\lim\limits_{s\to\infty} \widetilde{h}(s) = \infty$, and $\widetilde{h}(s) \le \frac{1}{2q}(qs + r) - \frac{r}{2q}\mathbb{1}_{\mathbb{R}_+}(r) \le \frac{s}{2}$ for all $s \in [s_1, \infty)$. The interval $(0, s_1)$ is not so 'important', one can choose any increasing $\widetilde{h}$ there, such that $\widetilde{h}(s) < \min\{\frac{s}{2}, \widetilde{h}(s_1)\}$, $s \in (0, s_1)$. By Proposition 2.16, (2.10) is equivalent to (2.3). Then, by (3.4), we have

$$\begin{aligned}\sup_{|\tau|\le \widetilde{h}(s)} \left| \frac{\widetilde{b}(s+\tau)}{\widetilde{b}(s)} - 1 \right| &= \sup_{q|\tau| \le h(qs+r) - \frac{r}{2}\mathbb{1}_{\mathbb{R}_+}(r)} \left| \frac{b(qs+r+q\tau)}{b(qs+r)} - 1 \right| \\ &\le \sup_{q|\tau|\le h(qs+r)} \left| \frac{b(qs+r+q\tau)}{b(qs+r)} - 1 \right| \to 0,\end{aligned}$$

as $s \to \infty$. Therefore, again by Proposition 2.16, (2.10) holds for b replaced by $\widetilde{b}$. Next, set $\nu(r) := \frac{r}{2}$, $r \ge 0$, and $\nu(r) := r$, $r < 0$, then

$$\begin{aligned}\widetilde{b}\big(\widetilde{h}(s)\big) s^{1+\delta} &= pb\big(h(qs+r) + \nu(r)\big) s^{1+\delta} \\ &= p \frac{b\big(h(qs+r) + \nu(r)\big)}{b\big(h(qs+r)\big)} b\big(h(qs+r)\big)(qs+r)^{1+\delta} \Big(\frac{s}{qs+r}\Big)^{1+\delta} \to 0,\end{aligned}$$

as $s \to \infty$, because of (2.1), (3.1). Therefore, $\widetilde{b} \in \mathcal{S}_{\rm reg}$. Finally, $b \in \mathcal{S}_{{\rm reg},n}$ for some $n \in \mathbb{N}$, trivially implies $\widetilde{b} \in \mathcal{S}_{{\rm reg},n}$.

(iii) Evidently, the convergence (2.10) implies the same one with b replaced by b_α, with the same h and for any $\alpha \in (0,1)$. Next, let $\alpha' \in (0,1)$ be such that $b_{\alpha'} \in L^1(\mathbb{R}_+)$. By the well-known log-convexity of L^p-norms (for $p > 0$), for any $\alpha \in (\alpha', 1)$ and for $\beta := \frac{\alpha - \alpha'}{\alpha(1-\alpha')} \in (0,1)$, we have $\frac{1}{\alpha} = \frac{1-\beta}{\alpha'} + \beta$ and

$$\|b\|_{L^\alpha(\mathbb{R}_+)} \le \|b\|^{1-\beta}_{L^{\alpha'}(\mathbb{R}_+)} \|b\|^{\beta}_{L^1(\mathbb{R}_+)} < \infty, \tag{3.7}$$

i.e. $b_\alpha \in L^1(\mathbb{R}_+)$ for all $\alpha \in (\alpha', 1)$. Take and fix now, an arbitrary $\alpha_0 \in \Big(\max\Big\{\alpha', \dfrac{1}{1+\delta}\Big\}, 1\Big)$. Then, for any $\alpha \in [\alpha_0, 1]$, we have that $\delta' := \alpha(1+\delta) - 1 \in (0, \delta]$, and hence, by (3.1),

$$\lim_{s\to\infty} b_\alpha\big(h(s)\big) s^{1+\delta'} = \lim_{s\to\infty} \Big(b\big(h(s)\big) s^{1+\delta}\Big)^\alpha = 0.$$

Therefore, $b_\alpha \in \mathcal{S}_{\rm reg}$, $\alpha \in [\alpha_0, 1]$.

Let, additionally, (3.2) hold for both b and $b_{\alpha'}$ (i.e., in particular, $b \in \mathcal{S}_{\mathrm{reg},n}$) and for some $n \in \mathbb{N}$. Then one can use again the log-convexity of L^p-norms, now for $L^p\big((\rho,\infty), s^n\,ds\big)$ spaces, to deduce that $b_\alpha \in \mathcal{S}_{\mathrm{reg},n}$, $\alpha \in [\alpha_0, 1]$.

Finally, $b, b_{\alpha_0} \in \mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$, implies $b, b_{\alpha_0} \in \mathcal{S}_{\mathrm{reg},1}$, and hence, cf. Remark 3.3, b and b_{α_0} are strongly sub-exponential on $\mathbb{R}$, i.e. (2.8) holds for both b and b_{α_0}. Therefore, for an arbitrary $\varepsilon \in (0,1)$, there exists $\rho_0 = \rho_0(\varepsilon, b, b_{\alpha_0}) > \rho$ (where ρ is from Definition 3.1) and

$$B_0 := 2(1+\varepsilon)\max\left\{\int_{\mathbb{R}} b(s)\,ds, \int_{\mathbb{R}} b_{\alpha_0}(s)\,ds\right\} > 0,$$

such that, for all $s \geq \rho_0$,

$$\begin{aligned} &\int_{\mathbb{R}} b(s-\tau)b(\tau)\,d\tau \leq B_0 b(s), \\ &\int_{\mathbb{R}} b_{\alpha_0}(s-\tau)b_{\alpha_0}(\tau)\,d\tau \leq B_0 b_{\alpha_0}(s). \end{aligned} \tag{3.8}$$

Then, applying again the norm log-convexity arguments, cf. (3.7), one gets, for any fixed $s \geq \rho_0$ and for all $\alpha \in (\alpha_0, 1)$

$$\int_{\mathbb{R}} \big(b(s-\tau)b(\tau)\big)^\alpha d\tau \leq \left(\int_{\mathbb{R}} \big(b(s-\tau)b(\tau)\big)^{\alpha_0} d\tau\right)^{\frac{1}{\alpha_0}(1-\beta)\alpha} \left(\int_{\mathbb{R}} b(s-\tau)b(\tau)d\tau\right)^{\beta\alpha},$$

where $\beta = \frac{\alpha-\alpha_0}{\alpha(1-\alpha_0)} \in (0,1)$. Combining the latter inequality with (3.8), one gets

$$\int_{\mathbb{R}} \big(b(s-\tau)b(\tau)\big)^\alpha d\tau \leq \big(B_0(b(s))^{\alpha_0}\big)^{\frac{1}{\alpha_0}(1-\beta)\alpha} \big(B_0 b(s)\big)^{\beta\alpha} = B_0\big(b(s)\big)^\alpha.$$

The theorem is fully proved now. □

It is naturally to expect that asymptotically small changes in the behavior at infinity preserves the sub-exponential property of a function. Namely, consider the following definition.

Definition 3.6. Two functions $b_1, b_2 : \mathbb{R} \to \mathbb{R}_+$ are said to be *weakly tail-equivalent* if

$$0 < \liminf_{s\to\infty} \frac{b_1(s)}{b_2(s)} \leq \limsup_{s\to\infty} \frac{b_1(s)}{b_2(s)} < \infty,$$

or, in other words, if there exist $\rho > 0$ and $C_2 \geq C_1 > 0$, such that,

$$C_1 b_1(s) \leq b_2(s) \leq C_2 b_1(s), \quad s \geq \rho. \tag{3.9}$$

Proposition 3.7. *Let $b_1 : \mathbb{R} \to \mathbb{R}_+$ be a weakly sub-exponential on $\mathbb{R}$ function. Let $b_2 : \mathbb{R} \to \mathbb{R}_+$ be a long-tailed function which is weakly tail-equivalent to b_1. Then b_2 is weakly sub-exponential on $\mathbb{R}$ as well. If, additionally,* (2.9) *holds for $b = b_1$, then b_2 is strongly sub-exponential on $\mathbb{R}$.*

Proof. Let b_1 be weakly sub-exponential on $\mathbb{R}$, cf. Definition 2.8, and the functions $b_{1,+}$ and $b_{2,+}$ be defined according to (2.4). Then, evidently, $b_{1,+}$ and $b_{2,+}$ will be also weakly tail-equivalent, and, moreover, $b_{2,+}$ will be long-tailed. Then, by [13, Theorem 4.8], $b_{2,+}$ is a also sub-exponential density on $\mathbb{R}_+$, i.e. (2.5) holds

for $b_+ = b_{2,+}$. As a result, by Definition 2.8, b_2 is weakly sub-exponential on $\mathbb{R}$. Next, let (2.9) holds for $b = b_1$. Then, by (3.9), we have, for all $s \geq \rho$,

$$b_2(s+\tau) \leq C_2 b_1(s+\tau) \leq C_2 K b_1(s) \leq \frac{C_2}{C_1} K b_2(s),$$

i.e. (2.9) holds for $b = b_2$ as well. As a result, by Lemma 2.13, both b_1 and b_2 are strongly sub-exponential on $\mathbb{R}$. □

Proposition 3.8. *Let $b_1 \in \mathcal{S}_{\mathrm{reg}}$ and let $b_2 : \mathbb{R} \to \mathbb{R}_+$ be a bounded tail-decreasing and tail-log-convex function, such that*

$$\lim_{s\to\infty} \frac{b_2(s)}{b_1(s)} = C \in (0,\infty). \tag{3.10}$$

Then $b_2 \in \mathcal{S}_{\mathrm{reg}}$.

Proof. First, we note that (3.10), (2.1) yield that b_2 is long-tailed as b_1 is such. Let $\delta \in (0,1)$ and $h : (0,\infty) \to (0,\infty)$ be an increasing function, such that $h(s) < \dfrac{s}{2}$, $\lim\limits_{s\to\infty} h(s) = \infty$, and (2.10) and (3.1) hold for $b = b_1$. Next, take an arbitrary $\varepsilon \in \big(0, \min\{1, C\}\big)$. Choose $\rho > 1$ such that b_2 is decreasing and log-convex on $[\rho,\infty)$, and $b_2(\rho) \leq 1$. By (3.10) and (2.10) (for $b = b_1$), there exists $\rho_1 \geq \rho$, such that, for all $s \geq \rho_1$,

$$0 < (C-\varepsilon) b_1(s) \leq b_2(s) \leq (C+\varepsilon) b_1(s), \tag{3.11}$$

$$\left| \frac{b_1(s \pm h(s))}{b_1(s)} - 1 \right| < \varepsilon. \tag{3.12}$$

Since b_2 is bounded and $b_1 \in L^1(\mathbb{R}_+)$, we have from (3.11) that $b_2 \in L^1(\mathbb{R}_+)$. Moreover, by (3.11), for any $s \geq 2\rho_1$,

$$\begin{aligned}
\frac{C-\varepsilon}{C+\varepsilon}\left(\frac{b_1(s \pm h(s))}{b_1(s)} - 1\right) + \frac{C-\varepsilon}{C+\varepsilon} &= \frac{(C-\varepsilon) b_1(s \pm h(s))}{(C+\varepsilon) b_1(s)} \\
&\leq \frac{b_2(s \pm h(s))}{b_2(s)} \\
&\leq \frac{(C+\varepsilon) b_1(s \pm h(s))}{(C-\varepsilon) b_1(s)} = \frac{C+\varepsilon}{C-\varepsilon}\left(\frac{b_1(s \pm h(s))}{b_1(s)} - 1\right) + \frac{C+\varepsilon}{C-\varepsilon},
\end{aligned}$$

and, therefore, by (3.12),

$$\left| \frac{b_2(s \pm h(s))}{b_2(s)} - 1 \right| < \max\left\{ \varepsilon \frac{C+\varepsilon}{C-\varepsilon} + \frac{C+\varepsilon}{C-\varepsilon} - 1,\ \varepsilon \frac{C-\varepsilon}{C+\varepsilon} + 1 - \frac{C-\varepsilon}{C+\varepsilon} \right\}.$$

Since the latter expression may be arbitrary small, by an appropriate choice of ε, one gets that (2.10) holds for $b = b_2$. Finally, (3.1) for $b = b_1$, and (3.10) imply that (3.1) holds for $b = b_2$ and the same δ and h. □

Remark 3.9. In the assumptions of the previous theorem, if, additionally, $b_1 \in \mathcal{S}_{\mathrm{reg},n}$ for some $n \in \mathbb{N}$, and b_2 is integrable on $(-\infty, -\rho_2)$ for some $\rho_2 > 0$, then $b_2 \in \mathcal{S}_{\mathrm{reg},n}$ (because of (3.11) and the boundedness of b_2).

On the other hand, if one can check that both functions b_1 and b_2 satisfy (2.10) with the same function $h(s)$, then the sufficient condition to verify (3.1) for $b = b_2$, provided that it holds for $b = b_1$, is much weaker than (3.10). To present the corresponding statement, consider the following definition.

Definition 3.10. Let $b_1, b_2 : \mathbb{R} \to \mathbb{R}_+$ and, for some $\rho \geq 0$, $b_i(s) > 0$ for all $s \in [\rho, \infty)$, $i = 1, 2$. The functions b_1 and b_2 are said to be *(asymptotically) log-equivalent*, if

$$\log b_1(s) \sim \log b_2(s), \quad s \to \infty. \tag{3.13}$$

Proposition 3.11. *Let $b_1 \in \mathcal{S}_{\mathrm{reg}}$ and let h be the function corresponding to Definition 3.1 with $b = b_1$. Let $b_2 : \mathbb{R} \to \mathbb{R}_+$ be a bounded tail-decreasing and tail-log-convex function, such that (2.10) holds with $b = b_2$ and the same h. Suppose that b_1 and b_2 are log-equivalent. Then $b_2 \in \mathcal{S}_{\mathrm{reg}}$. If, additionally, there exists $\alpha' \in (0, 1)$, such that (3.2) holds with $b = (b_1)^{\alpha'}$ and b_2 is integrable on $(-\infty, \rho)$, then $b_2 \in \mathcal{S}_{\mathrm{reg},n}$.*

Proof. Let $\delta \in (0, 1)$ be such that (3.1) holds for b replaced by b_1. Take an arbitrary $\varepsilon \in \big(0, \frac{\delta}{1+\delta}\big)$. By (3.13), there exists $\rho_\varepsilon > 0$, such that $b_i(s) < 1$, $s > \rho_\varepsilon$, $i = 1, 2$, and

$$\begin{gathered} -(1-\varepsilon)\log b_1(s) \leq -\log b_2(s) \leq -(1+\varepsilon)\log b_1(s), \quad s > \rho_\varepsilon, \\ b_1(s)^{1+\varepsilon} \leq b_2(s) \leq b_1(s)^{1-\varepsilon}, \quad s > \rho_\varepsilon. \end{gathered} \tag{3.14}$$

Since $h(s) \to \infty$, $s \to \infty$, there exists $\rho_0 > \rho_\varepsilon$, such that $h(s) > \rho_\varepsilon$ for any $s > \rho_0$. Then, by (3.14), we have, for all $s > \rho_0$,

$$b_2\big(h(s)\big)s^{(1+\delta)(1-\varepsilon)} < b_1\big(h(s)\big)^{1-\varepsilon} s^{(1+\delta)(1-\varepsilon)} = \big(b_1(h(s))s^{1+\delta}\big)^{1-\varepsilon},$$

and therefore, (3.1) holds with $b = b_2$ and δ replaced by

$$(1+\delta)(1-\varepsilon) - 1 = \delta - \varepsilon(1+\delta) \in (0, 1),$$

that proves the first statement. To prove the second one, assume, additionally, that $\varepsilon < 1 - \alpha'$. Then, by (3.14), we have, for all $s > \rho_\varepsilon$,

$$b_2(s)s^{n-1} \leq b_1(s)^{1-\varepsilon} s^{n-1} < b_1(s)^{\alpha'} s^{n-1},$$

as $b_1(s) < 1$ here. □

3.2 Examples

We consider now main examples of functions $b \in \mathcal{S}_{\mathrm{reg}}$. Because of Proposition 3.11, we will classify these functions 'up to log-equivalence', i.e. by the asymptotic behavior of

$$l(s) := -\log b(s).$$

Taking into account the result of Theorem 3.5 concerning the function $\breve{b}$, it will be enough to define b on some (s_0, ∞), $s_0 > 0$ only. Next, by Lemma 2.9, the function b_+ defined by (2.4) is a sub-exponential density on $\mathbb{R}_+$. Therefore, one can use the classical examples of such densities, see e.g. [13]. However, using the

result of Theorem 3.5 concerning the function $\widetilde{b}$, one can consider that examples in their 'simplest' forms (ignoring any shifts of the argument or scales of the argument or the function itself).

Now we consider different asymptotics of the function $l(s) = -\log b(s)$. In all particular examples below, it is straightforward to check that each particular bounded functions b is such that $b'(s) < 0$ and $(\log b(s))'' > 0$ for all big enough values of s, i.e. b is tail-decreasing and tail-log-convex.

Class 1: $l(s) \sim D \log s$, $s \to \infty$, $D > 1$

Polynomial decay Let $b : \mathbb{R} \to \mathbb{R}_+$ be a bounded tail-decreasing tail-log-convex function, such that

$$b(s) \sim q s^{-D}, \quad s \to \infty,\, D > 1,\; q > 0.$$

By Proposition 3.8, to show that $b \in \mathcal{S}_{\mathrm{reg}}$, it is enough to prove this for

$$b(s) = \mathbb{1}_{\mathbb{R}_+}(s) \frac{1}{(1+s)^D}, \quad s \in \mathbb{R}.$$

For an arbitrary $\gamma \in (0,1)$, consider $h(s) = s^\gamma$, $s > 0$. Then

$$\frac{b(s \pm h(s))}{b(s)} = \left(\frac{1+s}{1+s \pm s^\gamma} \right)^D \to 1, \quad s \to \infty.$$

Finally,

$$b\big(h(s)\big) s^{1+\delta} = \frac{s^{1+\delta}}{(1+s^\gamma)^D} \to 0, \quad s \to \infty,$$

provided that we will choose h above with $\gamma \in \big(\frac{1}{D}, 1\big)$ and take $\delta \in (0, \gamma D - 1) \subset (0,1)$. As a result, $b \in \mathcal{S}_{\mathrm{reg}}$. Clearly, $b \in \mathcal{S}_{\mathrm{reg},n}$ for $D > n$, cf. Remark 3.9. Consider now several classical examples.

Example 3.12. 1) *Student's t-function.* Let, for $p > \frac{1}{2}$,

$$\mathscr{T}(s) = \frac{1}{(1+s^2)^p}, \quad s > 0.$$

The probability density of Student's t-distribution is given, for $p = \frac{\nu+1}{2}$, $\nu > 0$, by $\frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\nu\pi}\Gamma(\frac{\nu}{2})} \mathscr{T}\big(\frac{s}{\sqrt{\nu}}\big)$, and extended symmetrically on the whole $\mathbb{R}$. Then $\mathscr{T} \in \mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$, if only $p > \frac{n}{2}$. The case $p = 1$ is referred to the Cauchy distribution, the corresponding function belongs to $\mathcal{S}_{\mathrm{reg},n}$ for $n = 1$ only.

2) *The Lévy function.* Let, for $c > 0$,

$$\mathscr{L}(s) = s^{-\frac{3}{2}} \exp\Big(-\frac{c}{s}\Big), \quad s > 0.$$

The probability density of the Lévy distribution is $\sqrt{\frac{c}{\pi}} \mathscr{L}(s - \mu)$, $\mu \in \mathbb{R}$, $s > \mu$.

3) *The Burr function.* Let, for $c > 0$, $k > 0$,

$$\mathscr{B}(s) = \frac{s^{c-1}}{(1+s^c)^{k+1}}, \quad s > 0.$$

The probability density of the so-called Burr IV distribution is just $ck\mathscr{B}(s)$. Note that the case $c = 1$ is related to the Pareto distribution; the latter has the density $kp^k\mathscr{B}(s-1)\mathbb{1}_{[p,\infty)}(s)$ for any $p > 0$.

Logarithmic perturbation of the polynomial decay Let $D > 1$, $\nu \in \mathbb{R}$, and

$$b(s) = \mathbb{1}_{(1,\infty)}(s)\frac{(\log s)^\nu}{s^D}, \quad s \in \mathbb{R}.$$

We are going to apply Proposition 3.11 now, with $b_1(s) = s^{-D}$ and $b_2(s) = (\log s)^\nu s^{-D}$. Indeed, then (3.13) evidently holds. It remains to check that (2.10) holds for both b_1 and b_2 with the same $h(s) = s^\gamma$, $\gamma \in (0,1)$. One has

$$\frac{\log(s \pm s^\gamma)}{\log s} = \frac{\log s + \log\bigl(1 \pm s^{\gamma-1}\bigr)}{\log s} \to 1, \quad s \to \infty,$$

that yields the needed.

Class 2: $l(s) \sim D(\log s)^q$, $s \to \infty$, $q > 1$, $D > 0$

Consider the function

$$N(s) := \mathbb{1}_{\mathbb{R}_+}(s)\exp\bigl(-D(\log s)^q\bigr), \quad s \in \mathbb{R}.$$

Take $h(s) = \mathbb{1}_{[\rho,\infty)}(s)s^{\frac{1}{q}}$, where $\rho > 1$ is chosen such that $h(s) < \frac{s}{2}$ for $s \geq \rho$. Prove that (2.10) holds. We have

$$\frac{N(s \pm h(s))}{N(s)} = \exp\biggl\{D(\log s)^q\biggl(1 - \Bigl(1 + \frac{\log(1 \pm s^{\frac{1}{q}-1})}{\log s}\Bigr)^q\biggr)\biggr\}.$$

Since $q > 1$, we have that $t(s) := \dfrac{\log(1 \pm s^{\frac{1}{q}-1})}{\log s} \to 0$, $s \to \infty$. Redefine then ρ to have that $|t(s)| < 1$, if only $s > \rho$. Use the binomial series

$$(1+t)^\alpha = \sum_{k=0}^{\infty}\binom{\alpha}{k}t^k, \quad \alpha > 0, \tag{3.15}$$

which converges for $|t| < 1$. One gets then (for $\alpha = q$)

$$\frac{N(s \pm h(s))}{N(s)} = \exp\biggl(-D(\log s)^q\sum_{k=1}^{\infty}\binom{q}{k}t(s)^k\biggr) \to 1, \quad s \to \infty.$$

Indeed, using the well-known inequality

$$\left|\binom{\alpha}{k}\right| \leq \frac{M}{k^{1+\alpha}}, \quad k \in \mathbb{N},\ M = M(\alpha), \tag{3.16}$$

one gets, for any $s > \rho$,

$$\left| -D(\log s)^q \sum_{k=1}^{\infty} \binom{q}{k} t(s)^k \right| \le D(\log s)^q |t(s)| \sum_{k=1}^{\infty} \frac{M}{k^{1+q}} \to 0, \quad s \to \infty,$$

since the latter series converges, and, for $q > 1$,

$$(\log s)^q |t(s)| \sim (\log s)^{q-1} s^{\frac{1}{q}-1} \to 0, \quad s \to \infty.$$

Finally, for any $\delta \in \mathbb{R}$,

$$N\big(s^{\frac{1}{q}}\big) s^{1+\delta} = \exp\big(-Dq^{-q}(\log s)^q + (1+\delta)\log s\big) \to 0, \quad s \to \infty,$$

since $q > 1$.

As a result, $N \in \mathcal{S}_{\mathrm{reg}}$. Moreover, evidently, $N \in \mathcal{S}_{\mathrm{reg},n}$, for any $n \in \mathbb{N}$.

We may also consider Proposition 3.11 for $b_1 = b$ and $b_2 = pb$, where b_2 is tail-decreasing and tail-log-convex function, such that $\log p = o(\log b)$ (that is equivalent to $\log b_1 \sim \log b_2$) and p satisfies (2.10) with $h(s) = s^{\frac{1}{q}}$. According to the results above, a natural example of such $p(s)$ might be s^D, $D \in \mathbb{R}$. It is straightforward to verify that, for any $D \in \mathbb{R}$, $b_2 = pb_1$ is tail-decreasing and tail-log-convex. As a result, then $b_2 \in \mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$. Consider now a classical example of such function b_2.

Example 3.13. *The log-normal function.* Let, for $\gamma > 0$,

$$\mathscr{N}(s) = \frac{1}{s} \exp\Big(-\frac{(\log s)^2}{2\gamma^2}\Big), \quad s > 0.$$

By the above, $\mathscr{N} \in \mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$. The log-normal distribution has the density $\frac{1}{\gamma\sqrt{2\pi}} \mathscr{N}(se^{-\mu})$ for an arbitrary $\mu \in \mathbb{R}$.

Class 3: $l(s) \sim s^{\alpha}$, $\alpha \in (0,1)$

Consider, for any $\alpha \in (0,1)$, the so-called *fractional exponent*

$$w(s) = \mathbb{1}_{\mathbb{R}_+}(s) e^{-s^{\alpha}}, \quad s \in \mathbb{R}. \tag{3.17}$$

Set $h(s) = \mathbb{1}_{[\rho,\infty)}(s)(\log s)^{\frac{2}{\alpha}}$, where $\rho > 0$ is chosen such that $h(s) < \frac{s}{2}$ for $s \ge \rho$. Then, in particular, $t(s) := \dfrac{h(s)}{s} < 1$, $s \ge \rho$. Prove that (2.10) holds. Using (3.15) for $t = \pm t(s)$, one gets

$$\frac{w\big(s \pm h(s)\big)}{w(s)} = \exp\bigg(-s^{\alpha} \sum_{k=1}^{\infty} \binom{\alpha}{k} (\pm t(s))^k\bigg) \to 1, \quad s \to \infty,$$

similarly to the arguments in Subsubsection 3.2, by using (3.16) and the evident convergence $s^{\alpha} t(s) \to 0$, $s \to \infty$, $\alpha \in (0,1)$. Finally, for any $\delta \in \mathbb{R}$,

$$w\big(h(s)\big) s^{1+\delta} = \exp\big(-(\log s)^2 + (1+\delta)\log s\big) \to 0, \quad s \to \infty.$$

As a result, $w \in \mathcal{S}_{\mathrm{reg}}$. It is clear also that $w \in \mathcal{S}_{\mathrm{reg},n}$ for all $n \in \mathbb{N}$. To find $\eta(t,w)$, one has to solve the equation $e^{-s^{\alpha}} = e^{-\beta t}$; therefore,

$$\eta(t,w) = (\beta t)^{\frac{1}{\alpha}}$$

Similarly to the above, one can show that $pw \in \mathcal{S}_{\mathrm{reg}}$, provided that, in particular, $\log p = o(\log w)$ and (2.10) holds for $b = p$ and $h(s) = (\log s)^{\frac{2}{\alpha}}$. Again, one can consider $p(s) = s^D$, $D \in \mathbb{R}$, since it satisfies (2.10) with $h(s) = s^\gamma > (\log s)^{\frac{2}{\alpha}}$, $\alpha, \gamma \in (0,1)$, and big enough s. As before, the verification that, for any $D \in \mathbb{R}$, $b_2 = pb_1$ is tail-decreasing and tail-log-convex is straightforward. Consider the corresponding classical example.

Example 3.14. *The Weibull function.* Let, for $\alpha \in (0,1)$,

$$\mathscr{W}(s) = \frac{\exp(-s^\alpha)}{s^{1-\alpha}}, \quad s \geq \rho > 0.$$

Note that $\int_s^\infty \mathscr{W}(\tau)\, d\tau = \frac{1}{\alpha} w(s)$, where w is given by (3.17). By the above, $\mathscr{W} \in \mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$. The probability density of the Weibull distribution is $\frac{\alpha}{\beta}\mathscr{W}\big(\frac{s}{\beta}\big)$, $s > 0$ for any $\beta > 0$. Note that the density itself is unbounded near 0.

Class 4: $l(s) \sim \dfrac{s}{(\log s)^\alpha}$, $\alpha > 1$

Consider also a function which decays 'slightly' slowly than an exponential function. Namely, let, for an arbitrary fixed $\alpha > 1$,

$$g(s) = \mathbb{1}_{\mathbb{R}_+}(s) \exp\Big(-\frac{s}{(\log s)^\alpha}\Big), \quad s \in \mathbb{R}.$$

Take, for an arbitrary $\gamma \in (1, \alpha)$, $h(s) = (\log s)^\gamma$, $s > 0$; and denote, for a brevity, $p(s) := \dfrac{h(s)}{s} \to 0$, $s \to \infty$. Then, $\log(s + h(s)) = \log s + \log(1 + p(s))$. Set also

$$q(s) = \frac{\log(1 + p(s))}{\log s} \to 0, \quad s \to \infty.$$

Then, for any $s > e^{\alpha+1}$, we have

$$\begin{aligned}
&\log \frac{g(s + h(s))}{g(s)} = \frac{s}{(\log s)^\alpha}\Big(1 - \frac{1 + p(s)}{\big(1 + q(s)\big)^\alpha}\Big) \\
&= \frac{1}{\big(1 + q(s)\big)^\alpha}\left(\alpha \frac{\big(1 + q(s)\big)^\alpha - 1}{\alpha q(s)} \frac{\log(1 + p(s))}{p(s)} (\log s)^{\gamma-\alpha-1} - (\log s)^{\gamma-\alpha}\right) \\
&\to 0, \quad s \to \infty,
\end{aligned}$$

as $\gamma < \alpha$; and similarly $\log \dfrac{g(s - h(s))}{g(s)} \to 0$, $s \to \infty$. Therefore, (2.10) holds for $b = g$. Next,

$$\log\big(g(h(s))s^{1+\delta}\big) = -(\log s)\Big(\frac{(\log s)^{\gamma-1}}{\gamma^\alpha (\log\log s)^\alpha} - (1 + \delta)\Big) \to -\infty, \quad s \to \infty,$$

that yields (3.1) for $b = g$. As a result, $g \in \mathcal{S}_{\mathrm{reg}}$. Again, evidently, $g \in \mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$. The same arguments as before show that, for any $D \in \mathbb{R}$, the function $s^D g(s)$ belongs to $\mathcal{S}_{\mathrm{reg},n}$ as well.

Remark 3.15. Naturally, $\alpha \in (0,1]$ gives behavior of $g(s)$ more 'close' to the exponential function. Unfortunately, our approach does not cover this case: the analysis above shows that $h(s)$, to fulfill even (2.6), must grow faster than $\log s$, whereas so 'big' $h(s)$ would not fulfill (2.10). In general, Lemma 2.9 gives a sufficient condition only, to get a sub-exponential density on $\mathbb{R}_+$. It can be shown, see e.g. [10, Example 1.4.3], that a probability distribution, whose density b on $\mathbb{R}_+$ is such that $\int_s^\infty b(\tau)\,d\tau \sim g(s)$, $s \to \infty$, with $\alpha > 0$, is a sub-exponential distribution (for the latter definition, see e.g. [13, Definition 3.1]). Then we expect that $b(s) \sim -g'(s)$, $s \to \infty$, and it is easy to see that $\log(-g'(s)) \sim \log g(s)$, $s \to \infty$. It should be stressed though that, in general, sub-exponential property of a distribution does not imply the corresponding property of its density, cf. [13, Section 4.2]. Therefore, we can not state that the function b above is a sub-exponential one for $\alpha \in (0,1]$.

Combining the results above, one gets the following statement.

Corollary 3.16. *Let $b : \mathbb{R} \to \mathbb{R}_+$ be a bounded tail-decreasing and tail-log-convex function, such that, for some $C > 0$, the function $Cb(s)$ has either of the following asymptotics as $s \to \infty$*

$$(\log s)^\mu s^{-(n+\delta)}, \qquad (\log s)^\mu s^\nu \exp\bigl(-D(\log s)^q\bigr),$$
$$(\log s)^\mu s^\nu \exp\bigl(-s^\alpha\bigr), \qquad (\log s)^\mu s^\nu \exp\Bigl(-\frac{s}{(\log s)^\gamma}\Bigr),$$

where $D, \delta > 0$, $q, \gamma > 1$, $\alpha \in (0,1)$, $\nu, \mu \in \mathbb{R}$. Then $b \in \mathcal{S}_{\mathrm{reg},n}$, $n \in \mathbb{N}$.

4 Analogues of Kesten's bound on $\mathbb{R}^d$

We start with a simple corollary of Propositions 2.20 and 2.21.

Proposition 4.1. *Let a function $\omega : \mathbb{R}^d \to (0, +\infty)$ be such that* (2.21) *holds, and, cf.* (2.22),

$$\limsup_{\lambda \to 0+} \sup_{x \in \Omega_\lambda} \frac{(a * \omega)(x)}{\omega(x)} \le 1. \tag{4.1}$$

Let also $a \in L^\infty(\mathbb{R}^d)$ and $\|a\|_\omega < \infty$. Then, for any $\delta \in (0,1)$, there exist $c_\delta > 0$ and $\lambda = \lambda(\delta) \in (0,1)$, such that

$$a^{*n}(x) \le c_\delta (1+\delta)^{n-1} \min\bigl\{\lambda, \omega(x)\bigr\}, \quad x \in \mathbb{R}^d. \tag{4.2}$$

Proof. Take any $\delta \in (0,1)$. By Proposition 2.21, there exists $\lambda = \lambda(\delta, \omega) \in (0,1)$, such that (2.20) holds, with ϕ given by (2.23) and $\gamma = 1 + \delta$. Denote $\|a\|_\infty := \|a\|_{L^\infty(\mathbb{R}^d)}$. We have

$$\frac{a(x)}{\omega_\lambda(x)} \le \frac{\|a\|_\infty}{\lambda} \mathbb{1}_{\mathbb{R}^d \setminus \Omega_\lambda}(x) + \frac{a(x)}{\omega(x)} \mathbb{1}_{\Omega_\lambda}(x) \le \frac{\|a\|_\infty}{\lambda} + \|a\|_\omega =: c_\delta < \infty, \tag{4.3}$$

and one can apply Proposition 2.20 that yields the statement. □

We are going to find now, for a given probability kernel a, an appropriate ω to validate (4.1).

Definition 4.2. Let $\mathcal{D}_d$ be the set of all bounded functions $b : \mathbb{R} \to (0, \infty)$, such that b is tail-decreasing (cf. Definition 2.15) and

$$\int_0^\infty b(s) s^{d-1}\, ds < \infty.$$

Let $\widetilde{\mathcal{D}}_d \subset \mathcal{D}_d$ denote the subset of all functions from $\mathcal{D}_d$ which are (strictly) decreasing to 0 on the whole $\mathbb{R}_+$.

Remark 4.3. It is easy to see that, if $b^{\alpha_0} \in \widetilde{\mathcal{D}}_d$ for some $\alpha_0 \in (0,1)$, then $b^\alpha \in \widetilde{\mathcal{D}}_d$ for all $\alpha \in [\alpha_0, 1]$.

Consider an analogue of long-tailed functions on $\mathbb{R}^d$.

Lemma 4.4. *Let $b \in \mathcal{D}_d$ be is a long-tailed function (cf. Definition 2.1), and $c(x) = b(|x|)$, $x \in \mathbb{R}^d$. Then, for any $r > 0$,*

$$\lim_{|x|\to\infty} \sup_{|y|\le r} \left| \frac{c(x+y)}{c(x)} - 1 \right| = 0. \tag{4.4}$$

Proof. Take arbitrary $r > 0$ and $|x| \ge r$. Let b be decreasing on (ρ, ∞) for some $\rho > 0$. Then, for any $|y| \le r$ and $|x| > \rho + r$, we get

$$b(|x| - r) \ge b(|x| - |y|) \ge b(|x+y|) \ge b(|x| + |y|) \ge b(|x| + r).$$

Therefore, for such values of x and y,

$$\left| \frac{c(x+y)}{c(x)} - 1 \right| = \left| \frac{b(|x+y|)}{b(|x|)} - 1 \right| \le \max\left\{ 1 - \frac{b(|x|+r)}{b(|x|)}, \frac{b(|x|-r)}{b(|x|)} - 1 \right\},$$

and hence (2.1) implies (4.4). □

We will assume in the sequel, that a is bounded by a radially symmetric function, namely:

$$\begin{aligned} &\text{There exists } b^+ \in \widetilde{\mathcal{D}}_d, \text{ such that} \\ &a(x) \le b^+(|x|), \text{ for a.a. } x \in \mathbb{R}^d. \end{aligned} \tag{4.5}$$

We start with the following sufficient condition.

Proposition 4.5. *Let* (4.5) *hold with $b^+ \in \widetilde{\mathcal{D}}_d$ which is log-equivalent, cf. Definition 3.10, to the function b, given by*

$$b(s) := \mathbb{1}_{\mathbb{R}_+}(s) \frac{M}{(1+s)^{d+\mu}}, \quad s \in \mathbb{R}, \tag{4.6}$$

for some $\mu, M > 0$. Then there exists $\alpha_0 \in (0,1)$, such that, for all $\alpha \in (\alpha_0, 1)$, the function $\omega(x) = b(|x|)^\alpha$, $x \in \mathbb{R}^d$, satisfies (4.1).

Proof. Set $\alpha_0 := \dfrac{d + \frac{\mu}{2}}{d+\mu} \in (0,1)$. Take arbitrary $\alpha \in (\alpha_0, 1)$ and $\epsilon \in (0, 1-\alpha)$. Take also an arbitrary $\delta \in (0,1)$, and define $h(s) = s^\delta$, $s > 0$. By (3.14), applied to $b_1 = b$ and $b_2 = b^+$, there exists $s_\delta > 2r$ such that, for all $s > s_\delta$,

$$h(s) < \frac{s}{2}, \qquad b^+(s) \le \big(b(s)\big)^{1-\epsilon}. \tag{4.7}$$

For an arbitrary $x \in \mathbb{R}^d$ with $|x| > s_\delta$, we have a disjoint expansion $\mathbb{R}^d = D_1(x) \sqcup D_2(x) \sqcup D_3(x)$, where

$$D_1(x) := \big\{|y| \le h(|x|)\big\}, \quad D_2(x) := \Big\{h(|x|) < |y| \le \frac{|x|}{2}\Big\},$$
$$D_3(x) = \Big\{|y| \ge \frac{|x|}{2}\Big\}.$$

Then, $\frac{(a*\omega)(x)}{\omega(x)} = I_1(x) + I_2(x) + I_3(x)$, where

$$I_j(x) := \int_{D_j(x)} a(y) \frac{(1+|x|)^{(d+\mu)\alpha}}{(1+|x-y|)^{(d+\mu)\alpha}} dy, \quad j = 1, 2, 3.$$

Using the inequality $|x-y| \ge \big||x| - |y|\big|$, $x, y \in \mathbb{R}^d$, one has that $|x-y| \ge |x| - |y| \ge |x| - |x|^\delta$ for $y \in D_1(x)$, $|x| > s_\delta$. Then

$$I_1(x) \le \left(\frac{1+|x|}{1+|x|-|x|^\delta}\right)^{(d+\mu)\alpha} \int_{D_1(x)} a(y) dy \to 1, \quad |x| \to \infty.$$

Next, we evidently have, for any $|y| < \frac{|x|}{2}$, that $1 + |x-y| \ge 1 + |x| - |y| \ge \frac{1}{2}(1+|x|)$; therefore,

$$I_2(x) \le 2^{(d+\mu)\alpha} \int_{\{|y| > |x|^\delta\}} a(y) dy \to 0, \quad |x| \to \infty.$$

Finally, by (4.5) and (4.7), the inclusions $y \in D_3(x)$ and $|x| > s_\delta$ imply

$$a(y) \le b^+(|y|) \le b(|y|)^{1-\epsilon} \le b\Big(\frac{|x|}{2}\Big)^{1-\epsilon},$$

and, therefore,

$$\begin{aligned} I_3(x) &\le M \frac{(1+|x|)^{(d+\mu)\alpha}}{\Big(1+\frac{|x|}{2}\Big)^{(d+\mu)(1-\epsilon)}} \int_{D_3(x)} \frac{1}{(1+|x-y|)^{(d+\mu)\alpha_0}} dy \\ &\le M \frac{(1+|x|)^{(d+\mu)\alpha}}{\Big(1+\frac{|x|}{2}\Big)^{(d+\mu)(1-\epsilon)}} \int_{\mathbb{R}^d} \frac{1}{(1+|y|)^{d+\frac{\mu}{2}}} dy \to 0, \quad |x| \to \infty, \end{aligned}$$

as $1 - \epsilon > \alpha$. Since b is decreasing on $\mathbb{R}_+$, we have, by (2.21), that, for any $\lambda > 0$, there exists $\rho_\lambda > 0$, such that $\Omega_\lambda = \{x \in \mathbb{R}^d : |x| > \rho_\lambda\}$. As a result, one gets (4.1) from the above. □

Lemma 4.6. *Let $b \in L^1(\mathbb{R})$ be even, positive, decreasing to 0 on the whole $\mathbb{R}_+$, and long-tailed function. Suppose that there exist $B, r_b, \rho_b > 0$, such that*

$$\int_{r_b}^{\infty} b(s-\tau) b(\tau)\, d\tau \le B b(s), \quad s > \rho_b. \tag{4.8}$$

Suppose also that

$$\lim_{|x|\to\infty} \frac{a(x)|x|^{d-1}}{b(|x|)} = 0. \tag{4.9}$$

Then the inequality (4.1) *holds for $\omega(x) := b(|x|)$, $x \in \mathbb{R}^d$.*

Proof. The assumption (4.9) implies that

$$g(r) := \sup_{|x|\ge r} \frac{a(x)|x|^{d-1}}{\omega(x)} \to 0, \quad r\to\infty. \tag{4.10}$$

Take an arbitrary $\delta\in(0,1)$. By (4.10), one can take then $r=r(\delta)>r_b$ such that $g(r)<\delta$.

Next, by Lemma 4.4, the inequality (4.4) holds for $c=\omega$. Therefore, there exists $\rho=\rho(\delta,r)=\rho(\delta)>\max\{r,\rho_b\}$, such that

$$\sup_{|y|\le r} \frac{\omega(x-y)}{\omega(x)} < 1+\delta, \quad |x|\ge\rho. \tag{4.11}$$

Then, by (4.10) and (4.11), we have

$$\begin{aligned}(a*\omega)(x) &= \omega(x)\int_{\{|y|\le r\}} a(y)\frac{\omega(x-y)}{\omega(x)}dy \\ &\quad + \omega(x)\int_{\{|y|\ge r\}} \frac{a(y)|y|^{d-1}}{\omega(y)}\,\frac{\omega(x-y)\omega(y)}{\omega(x)|y|^{d-1}}dy \\ &\le \omega(x)(1+\delta)\int_{|y|\le r} a(y)dy \\ &\quad + g(r)\omega(x)\int_{\{|y|\ge r\}} \frac{b(|x-y|)b(|y|)}{b(|x|)|y|^{d-1}}dy,\end{aligned}$$

and using that b is decreasing on $\mathbb{R}_+$ and the inequality $|x-y|\ge\big||x|-|y|\big|$, one gets, cf. (2.19) and recall that $g(r)<\delta$,

$$\le \omega(x)(1+\delta)+\delta\omega(x)\int_{\{|y|\ge r\}} \frac{b\big(\big||x|-|y|\big|\big)b(|y|)}{b(|x|)|y|^{d-1}}dy,$$

and using the spherical coordinates, one gets

$$\le \omega(x)(1+\delta)+\delta\omega(x)\sigma_d\int_r^\infty \frac{b\big(|x|-p\big)b(p)}{b(|x|)}dp, \tag{4.12}$$

where σ_d is the hyper-surface area of a unit sphere in $\mathbb{R}^d$ (note that we have omitted an absolute value, as b is even). Finally, using that $r>r_b$ and $\rho>\rho_b$, we obtain from (4.8) and (4.12) that, for any $\delta\in(0,1)$,

$$(a*\omega)(x)\le\omega(x)\big(1+\delta(1+\sigma_d B)\big), \quad |x|>\rho(\delta),$$

that implies the statement. □

Lemma 4.7. *Let $b\in\mathcal{S}_{\mathrm{reg},1}$ be an even function. Suppose that there exists $\alpha'\in(0,1)$ such that $b^{\alpha'}\in\widetilde{\mathcal{D}}_d$, i.e.*

$$\int_0^\infty b(s)^{\alpha'}s^{d-1}\,ds<\infty, \tag{4.13}$$

and, for any $\alpha\in(\alpha',1)$,

$$\lim_{|x|\to\infty}\frac{a(x)}{b(|x|)^\alpha}|x|^{d-1}=0. \tag{4.14}$$

Then there exists $\alpha_0\in(\alpha',1)$ such that the inequality (4.1) holds for $\omega(x)=b(|x|)^\alpha$, $x\in\mathbb{R}^d$ for all $\alpha\in(\alpha_0,1)$.

Proof. We apply the second part of Theorem 3.5, for $n = 1$; note that then (4.13) implies (3.2). As a result, for any $\alpha \in (\alpha_0, 1)$, the inequality (3.6) holds; in particular, then (4.8) holds with b replaced by b^α. The latter together with (4.14) allows to apply Lemma 4.6 for b replaced by b^α, that fulfils the statement. □

Remark 4.8. Note that, by Remark 4.3, (4.13) implies that $b \in \widetilde{\mathcal{D}}_d$ and hence, cf. Definition 3.1, $b \in \mathcal{S}_{\mathrm{reg},d}$.

As a result, one gets a counterpart of Proposition 4.5, for the case when the function b^+ in (4.5) decays faster than polynomial and $d > 1$.

Proposition 4.9. *Let* (4.5) *hold for a function* $b^+ \in \widetilde{\mathcal{D}}_d$ *which is loq-equivalent to a function* $b \in \mathcal{S}_{\mathrm{reg},1}$. *For* $d > 1$, *we suppose, additionally, that*

$$\lim_{s\to\infty} b(s)s^\nu = 0, \qquad \text{for all } \nu \geq 1. \tag{4.15}$$

Then there exists $\alpha_0 \in (0, 1)$, *such that, for all* $\alpha \in (\alpha_0, 1)$, *the function* $\omega(x) = b(|x|)^\alpha$, $x \in \mathbb{R}^d$ *satisfies* (4.1).

Proof. We will use Lemma 4.7. For $d > 1$, one gets from (4.15) that, for any $\nu > 0$, there exists $\rho_\nu \geq 1$, such that $b(s) \leq s^{-\nu}$, $s > \rho_\nu$. In particular, for any $\alpha' \in (0, 1)$, one has (4.13). For $d = 1$, $\sigma = 0$, we use instead that $b \in \mathcal{S}_{\mathrm{reg},1}$ implies (3.3), and hence we get (4.13), if only $\alpha' \in \big(\frac{1}{1+\delta}, 1\big)$.

Next, for any $d \in \mathbb{N}$, choose an arbitrary $\alpha \in (\alpha', 1)$. Then, by (4.5) and (3.14) applied for $b_1 = b$ and $b_2 = b^+$, we have that, for any $\epsilon \in (0, 1-\alpha)$, there exists $\rho_\epsilon > 0$, such that, for all $|x| > \rho_\varepsilon$,

$$\frac{a(x)}{b(|x|)^\alpha}|x|^{d-1} \leq b(|x|)^{1-\epsilon-\alpha}|x|^{d-1} = \Big(b(|x|)|x|^\nu\Big)^{1-\epsilon-\alpha}, \tag{4.16}$$

where $\nu = \dfrac{d-1}{1-\epsilon-\alpha} \geq 0$, as $\alpha < 1 - \epsilon$. Clearly, (4.16) together with (4.15), in the case $d > 1$, imply (4.14), that fulfills the statement. □

Remark 4.10. Note that, in Proposition 4.5, for the function b given by (4.6), one can choose $\alpha' \in (0, 1)$ such that (4.13) holds. The same property we have checked above for the function b which satisfies assumptions of Proposition 4.9. As a result, by Remark 4.3, the functions $\omega(x) = b(|x|)^\alpha$, $x \in \mathbb{R}^d$ in these Propositions are integrable for all $\alpha \in (\alpha_0, 1)$.

Definition 4.11. Let the set $\widetilde{\mathcal{S}}_{\mathrm{reg},d} \subset \mathcal{S}_{\mathrm{reg},d}$, $d \in \mathbb{N}$ be defined as follows. Let $\widetilde{\mathcal{S}}_{\mathrm{reg},1}$ be just the set $\mathcal{S}_{\mathrm{reg},1}$, whereas, for $d > 1$, let $\widetilde{\mathcal{S}}_{\mathrm{reg},d}$ be the set of all functions $b \in \mathcal{S}_{\mathrm{reg},d}$, such that b is either given by (4.6) for some $M, \mu > 0$ or b satisfies (4.15).

Remark 4.12. All functions in Classes 2–4 in Subsection 3.2 evidently satisfy (4.15) and hence belong to $\widetilde{\mathcal{S}}_{\mathrm{reg},d}$.

Theorem 4.13. *Let* (4.5) *hold with* $b^+ \in \widetilde{\mathcal{D}}_d$ *which is log-equivalent to a function* $b \in \widetilde{\mathcal{S}}_{\mathrm{reg},d}$. *Then there exists* $\alpha_0 \in (0, 1)$, *such that, for any* $\delta \in (0, 1)$ *and* $\alpha \in (\alpha_0, 1)$, *there exist* $c_1 = c_1(\delta, \alpha) > 0$ *and* $\lambda = \lambda(\delta, \alpha) \in (0, 1)$, *such that*

$$a^{*n}(x) \leq c_1(1+\delta)^n \min\big\{\lambda, b(|x|)^\alpha\big\}, \quad x \in \mathbb{R}^d.$$

In particular, for some $c_2 = c_2(\delta, \alpha) > 0$ *and* $s_\alpha = s_\alpha(\delta) > 0$,

$$a^{*n}(x) \leq c_2(1+\delta)^n b(|x|)^\alpha, \quad |x| > s_\alpha, n \in \mathbb{N}. \tag{4.17}$$

Proof. Combining Proposition 4.5 and 4.9 with Proposition 4.1, we get by (4.2) and (4.3) that there exist $\tilde{c_\delta} = \tilde{c_\delta}(\omega)$, $\lambda = \lambda(\delta, \alpha) \in (0,1)$, where $\omega(x) = b(|x|)^\alpha$, $x \in \mathbb{R}^d$, such that

$$a^{*n}(x) \le \tilde{c}_\delta (1+\delta)^{n-1} \min\{\lambda, b(|x|)^\alpha\}, \quad x \in \mathbb{R}^d, \tag{4.18}$$

that evidently yields (4.18). Since b is tail-decreasing, we have that, for some $s_\alpha > 0$, $b(|x|)^\alpha < \lambda$ for $|x| > s_\alpha$. This implies (4.17). □

Corollary 4.14. *Let $a(x) = b(|x|)$, $x \in \mathbb{R}$ for some $b \in \widetilde{\mathcal{S}}_{\mathrm{reg},d}$. Then there exists $\alpha_0 \in (0,1)$, such that, for any $\delta \in (0,1)$ and $\alpha \in (\alpha_0, 1)$, there exist $c_{\delta,\alpha} > 0$ and $s_\alpha = s_\alpha(\delta) > 0$, such that, for all ,*

$$a^{*n}(x) \le c_{\delta,\alpha}(1+\delta)^n a(x)^\alpha, \quad |x| > s_\alpha, n \in \mathbb{N}.$$

Proof. Since, for some $\rho > 0$, b is decreasing on (ρ, ∞) and (3.2) holds, there exists $b^+ \in \widetilde{\mathcal{D}}_d$, such that $b^+(s) = b(s)$, $s > \rho$ and $b^+(s) \ge b(s)$, $s \in [0, \rho]$. Then one can apply Theorem 4.13. □

Appendix: Non-local heat equation

We can apply the obtained results to the study of the regular part of the fundamental solution to the non-local heat equation

$$\frac{\partial}{\partial t} u(x,t) = \varkappa \int_{\mathbb{R}^d} a(x-y)\big(u(y,t) - u(x,t)\big)\, dy, \quad x \in \mathbb{R}^d, \tag{A.1}$$

where $\varkappa > 0$ and $0 \le a \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ is normalized, i.e. $\int_{\mathbb{R}^d} a(x)\, dx = 1$; see e.g. [1, 5, 15]. Consider an initial condition $u(x,0) = u_0(x)$, $x \in \mathbb{R}^d$ to (A.1) with u_0 from a space E of bounded on $\mathbb{R}^d$ functions. Since the operator $Au = \varkappa a * u - \varkappa u$ in the right hand side of (A.1) is bounded on E, the unique solution to (A.1) is given by

$$u(x,t) = e^{-\varkappa t}\big((\delta_0 + \phi_\varkappa(t)) * u_0\big)(x), \tag{A.2}$$

where δ_0 is the Dirac delta at $0 \in \mathbb{R}^d$ and

$$\phi_\varkappa(x,t) := \sum_{n=1}^{\infty} \frac{\varkappa^n t^n}{n!} a^{*n}(x), \quad x \in \mathbb{R}^d, t \ge 0. \tag{A.3}$$

Note that it was shown in [6, Lemma 2.2], that if a is a rapidly decreasing smooth function, then ϕ_k is indeed the solution to (A.1) with $u_0 = \delta_0$.

Now, for $d = 1$, suppose that $a(x) = b(x)$, $x \in \mathbb{R}^1$, and b satisfies the conditions of Theorem 2.22. Then, by (2.27), the series in (A.3) converges uniformly on finite time intervals for each $x > s_0$, and therefore, by (2.18),

$$\phi_\varkappa(x,t) \sim kte^{\varkappa t} a(x), \quad x \to \infty,\ t > 0.$$

For $d > 1$, let a and b satisfy the conditions of Theorem 4.13. Then, for each $\delta > 0$ and for each $\alpha < 1$ close enough to 1,

$$\phi_\varkappa(x,t) \le c_{\delta,\alpha}\big(e^{\varkappa t(1+\delta)} - 1\big) b(|x|)^\alpha, \quad |x| > s_\alpha,\ t > 0 \tag{A.4}$$

for some $c_{\delta,\alpha} > 0$ and $s_\alpha = s_\alpha(\delta) > 0$. In particular, if a is radially symmetric and the conditions of Corollary 4.14 hold, then one can replace $b(|x|)$ on $a(x)$ in (A.4).

Moreover, combining (4.18) with (A.2), one can get an estimate for the solution u to (A.1) as well. The further analysis of solutions to (A.1) can be found in [11].

Acknowledgements

Financial supports by the DFG through CRC 701, Research Group "Stochastic Dynamics: Mathematical Theory and Applications", and by the European Commission under the project STREVCOMS PIRSES-2013-612669 are gratefully acknowledged.

References

[1] F. Andreu-Vaillo, J. M. Mazón, J. D. Rossi, and J. J. Toledo-Melero. *Nonlocal diffusion problems*, volume 165 of *Mathematical Surveys and Monographs*. AMS Providence, RI, 2010.

[2] S. Asmussen, S. Foss, and D. Korshunov. Asymptotics for sums of random variables with local subexponential behaviour. *J. Theoret. Probab.*, 16(2):489–518, 2003.

[3] K. B. Athreya and P. E. Ney. *Branching processes*. Springer-Verlag, New York-Heidelberg, 1972. Die Grundlehren der mathematischen Wissenschaften, Band 196.

[4] A. A. Borovkov and K. A. Borovkov. *Asymptotic analysis of random walks*, volume 118 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2008. Heavy-tailed distributions, Translated from the Russian by O. B. Borovkova.

[5] C. Brändle, E. Chasseigne, and R. Ferreira. Unbounded solutions of the nonlocal heat equation. *Communications on Pure and Applied Analysis*, 10(6):1663–1686, 2011.

[6] E. Chasseigne, M. Chaves, and J. D. Rossi. Asymptotic behavior for nonlocal diffusion equations. *Journal de Mathématiques Pures et Appliquées*, 86(3):271–291, 2006.

[7] V. P. Chistyakov. A theorem on sums of independent positive random variables and its applications to branching random processes. *Theor. Probab. Appl.*, 9:640–648, 1964.

[8] J. Chover, P. Ney, and S. Wainger. Functions of probability measures. *J. Analyse Math.*, 26:255–302, 1973.

[9] D. B. H. Cline and S. I. Resnick. Multivariate subexponential distributions. *Stochastic Process. Appl.*, 42(1):49–72, 1992.

[10] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling extremal events*, volume 33 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1997. For insurance and finance.

[11] D. Finkelshtein, Y. Kondratiev, and P. Tkachov. Accelerated front propagation for monostable equations with nonlocal diffusion. arXiv:1611.09329.

[12] S. Foss and D. Korshunov. Lower limits and equivalences for convolution tails. *Ann. Probab.*, 35(1):366–383, 2007.

[13] S. Foss, D. Korshunov, and S. Zachary. *An introduction to heavy-tailed and subexponential distributions*. Springer Series in Operations Research and Financial Engineering. Springer, New York, second edition, 2013.

[14] C. Klüppelberg. Subexponential distributions and characterizations of related classes. *Probab. Theory Related Fields*, 82(2):259–269, 1989.

[15] Y. Kondratiev, S. Molchanov, A. Piatnitski, and E. Zhizhina. Resolvent bounds for jump generators. *Applicable Analysis*, 2016. doi: 10.1080/00036811.2016.1263838.

[16] E. A. M. Omey. Subexponential distribution functions in $\mathbb{R}^d$. *J. Math. Sci. (N. Y.)*, 138(1):5434–5449, 2006.

[17] B. A. Rogozin and M. S. Sgibnev. Strongly subexponential distributions, and Banach algebras of measures. *Sibirsk. Mat. Zh.*, 40(5):1137–1146, 1999.

[18] G. Samorodnitsky and J. Sun. Multivariate subexponential distributions and their applications. *Extremes*, 19(2):171–196, 2016.

[19] M. S. Sgibnev. Banach algebras of functions with the same asymptotic behavior at infinity. *Siberian Math. J.*, 22(3):467–473, 1982.

[20] M. S. Sgibnev. The asymptotics of infinitely divisible distributions in $\mathbb{R}$. *Sibirsk. Mat. Zh.*, 31(1):135–140, 221, 1990.

[21] T. Watanabe. Convolution equivalence and distributions of random sums. *Probab. Theory Related Fields*, 142(3-4):367–397, 2008.